

Molecular Basis of Thyroid Cancer*

NADIR R. FARID, YUFEI SHI, AND MINJING ZOU

Molecular Endocrinology Laboratory, and Department of Medicine, King Faisal Specialist Hospital and Research Centre, Riyadh 11211, Kingdom of Saudi Arabia

- I. Introduction
- II. Thyroid Cell Growth
- III. Epidemiological and Clinical Considerations
- IV. Genetic Factors
- V. The Cell Cycle and Cancer
- VI. Thyroid Cell Cancer—Mono or Polyclonal?
- VII. Growth Factors and Their Receptors in Thyroid Tumors
- VIII. Oncogenes Involved in Thyroid Carcinogenesis
 - A. *ras*
 - B. *c-myc* and *c-fos*
 - C. PTC/RET oncogene
 - D. TRK-T1
 - E. *Met*
 - F. *Gsa* mutations
- IX. The TSH Receptor
- X. Tumor Suppressor Genes
 - A. LOH in thyroid tumors
 - B. Retinoblastoma (Rb) gene
 - C. p53
- XI. Progression to Metastatic Potential
- XII. nm23
- XIII. Perspective

Nature is nowhere accustomed more openly to display her secret mysteries than in cases where she shows traces of her workings apart from the beaten path; nor is there any better way to advance the proper practice of medicine than to give our minds to the discovery of the usual law of Nature by careful investigation of rarer forms of disease. For it has been found in almost all things, that what they contain of useful or applicable nature is hardly perceived unless we are deprived of them, or they become deranged in some way.

William Harvey (1657) (1)

I. Introduction

THYROID follicular cell tumors present a unique model for the study of genetic and environmental factors pre-

Address requests for reprints to: Nadir R. Farid, MBBS, FRCP, FRCP(C), FACP, Executive Director, Medical Research Institute, King Fahad National Guard Hospital, PO Box 22490, Riyadh 11426, Kingdom of Saudi Arabia.

*Supported by Grants 90-0014, 90-0015, and 90-0016 from King Faisal Specialist Hospital and Research Centre; Grant AT-13-2 from King Abdul Aziz City for Science and Technology (KACST); the Riyadh Chamber of Commerce; and Bristol-Myers-Squibb (Canada).

disposing to benign nodule formation, well differentiated malignant tumors, and anaplastic cancer. Although these histological changes are not necessarily sequential, there is evidence that gradation of proliferative and differentiative potential exists among cells in each thyroid follicle. Conditions conducive to rapid growth ensure that the progeny of cells with high growth potential establish local dominance, a prelude to nodule formation (2). There is also evidence that well differentiated carcinoma may progress to an anaplastic form (3). It is only speculation that benign nodules may develop into well differentiated carcinoma.

Cancer is a complex, multistep process (4). Based on the measurements of age-dependent tumor incidence, it was inferred mathematically that a succession of five or six independent steps are involved, each of which is rate limiting (5). This predicted sequence of events has been shown to apply to colorectal carcinoma (6-12). In the intact host, each step would represent a physiological barrier to be breached for a cell to progress to malignant transformation. The fact that multiple barriers must be overcome ensures that malignancy is a rare event (13).

A consideration of thyroid carcinogenesis raises several issues: What are the cellular events for the transformation of normal thyroid epithelial cells to malignant ones? Are the events that direct transformation of normal thyroid cells to follicular carcinoma and papillary carcinoma separate and independent? How can one account for tumors of mixed histology? How does iodide deficiency enhance the prevalence of follicular carcinoma? Are additional cellular events necessary for the metastatic potential of tumors or is this inherent in those changes resulting in follicular or papillary histology? Similarly, are additional factors involved in the progression of well differentiated carcinoma to anaplastic histology? Is there evidence that benign nodular proliferation is an antecedent to malignant transformation of thyroid cells as has been observed in colorectal cancer?

In this review, an attempt will be made to summarize what is known at the molecular level about thyroid cell growth and the factors involved in the transformation of thyroid cells. The picture is incomplete since what is known about thyroid tumors is not as advanced as for other tumor systems.

II. Thyroid Cell Growth

The thyroid follicular cell is a highly differentiated cell having evolved a highly efficient system for concentrating iodide for the synthesis, storage, and release of a reservoir of thyroid hormones, under basal conditions as well as condi-

tions of increased need. Thyroid cells are programmed (14) to aggregate into follicles (15) with a central space for the storage of thyroglobulin. This organization necessitates the polarization of the thyroid cell: the basal pole specializes in the import of iodide, which is then transported to the apex for iodination of thyroglobulin (16).

Unlike many other highly specialized cells, thyroid cells are not irreversibly terminally differentiated. When they proliferate in response to certain growth signals they temporarily lose the ability to concentrate iodide and to synthesize thyroglobulin (see Ref. 17 for review). Certain unregulated growth signals associated with malignant transformation may contribute to the loss of differentiated thyroid function. The thyroid cell turns over, on average, no more than five times through adulthood (18). Turnover increases in early infancy and adolescence (18, 19). Because of the inhomogeneity of the growth potential of follicular cells (2), a fraction of these follicular cells in adults would likely turnover much more than five times. It is assumed that cycles of cell division may be more frequent in benign or malignant tumors (2) although evidence from human tumors does not always support this notion (20).

An understanding of the factors involved in normal human thyroid cell growth, differentiation, and signaling is an essential prelude to understanding aberrations related to thyroid cell transformation. Unfortunately, the picture is incomplete due to the variability among species of factors that sustain thyroid proliferation, the pathways used, and cytokines elaborated. Data from animal thyroid tissue in primary culture or from thyroid rat cell lines may not be generalizable to the human thyroid (21-23) (see Ref. 16 for review).

TSH induces human thyroid cell growth at a higher concentration than is necessary for induction of differentiated function (24, 25). TSH mediates its growth stimulation through the adenyl cyclase/cAMP pathway. Insulin or insulin-like growth factor I (IGF-I) synergizes with TSH to induce thyroid cell growth while maintaining specialized cell function (25, 26). Insulin and IGF-I are thus either permissive for other factors without being mitogenic themselves or do not inhibit differentiated function (25, 26). Indeed, TSH has been shown to enhance insulin-induced autophosphorylation of both insulin and IGF-I receptors in rat thyroid cell lines as well as the phosphorylation of the immediate insulin receptor 185 kilodalton (kDa) endogenous substrate (27, 28).

Epidermal growth factor (EGF), on the other hand, induces human thyroid cell growth at the expense of loss in differentiated function (29-31). In some species, TSH induces EGF receptors on thyroid cells, making these cells more responsive to EGF (32, 33). Many EGF effects are reproduced by phorbol esters, used as probes for protein kinase C and diacylglycerol (DAG) (34-36). The effects of TSH, EGF, and phorbol esters on differentiation are largely independent of their mitogenic effects (17). Fibroblast growth factor (FGF) is also a potent thyroid cell mitogen (37).

Growth factors elaborated by thyrocytes have been studied in different species at different times. Human and sheep thyroids produce IGF-I (35, 38), FRTL5 cells produce IGF-II

(39), and porcine cells produce FGFs (40); both IGF-I and FGF act upon thyrocytes in an autocrine fashion.

Likewise, cytokines traditionally thought to be synthesized and secreted by immunocytes are synthesized, secreted, and act upon thyrocytes. These cytokines include interleukin-1 (IL-1), transforming growth factor β (TGF β), and IL-8. They either have growth-promoting effects as in physiological concentrations of IL-1 and IL-8 (41-43) or inhibit growth and in some instances differentiated function, e.g. TGF β , interferon- γ , and IL-1 (44-47). Indeed, TGF β may have a physiological role in negatively regulating thyroid follicular cell function. IL-1 has been shown paradoxically to limit the growth of human thyroid tumor cell lines (48). Several of these cytokines elaborated by thyrocytes synergize with each other and with growth factors (42, 45) in their effect on thyroid cell growth and differentiated function.

Many of the factors elaborated by the thyroid, e.g. IGF-I (38) and FGF (40) also act *in vivo* on endothelial cells and fibroblasts to induce neurovascularization (49) and to enhance adhesion molecule synthesis to support thyroid growth.

At least three distinct pathways for signal transduction have been defined in the thyroid: 1) receptor/adenylate cyclase/protein kinase A system; 2) receptor/tyrosine kinase pathway; and 3) receptor/phospholipase C cascade (50, 51). TSH activates both the adenylate cyclase and phospholipase C pathways. Activation of phospholipase C results in the formation of DAG and inositol-1,4,5-triphosphate (IP₃) (52, 53). DAG activates protein kinase C and IP₃ increases intracellular calcium concentrations. The concentration of TSH necessary to activate phospholipase C is much higher and its action slower than that necessary for adenylate cyclase activation. Apparently cAMP induces DAG synthetase in the thyroid, thus providing substrate for protein kinase C, allowing the activation of this pathway at physiological TSH concentrations (54).

EGF, insulin, and IGF-I act through tyrosine protein kinase receptors (17). EGF also mobilizes calcium (Ca⁺⁺) from intracellular stores as well as from extracellular sources through the generation of IP₃, at least in some species (17, 55, 56), and perhaps through other mechanisms.

A role for iodide in the regulation of thyroid cell growth has been proposed. Apparently through oxidized intermediaries, iodide may decrease both adenylate cyclase and Ca⁺⁺ levels in thyroid cells, thus reducing their sensitivity to TSH signaling (57-61). Iodide deficiency, conversely, enhances the effects of TSH on the thyroid.

TSH induces or increases phosphorylation of at least 11 proteins. EGF induces the phosphorylation of five proteins including mitogen-activated protein kinase, which in turn phosphorylates ribosomal S6 kinase (pp90^{rk}) (62-64). Phorbol esters induce the phosphorylation of 19 proteins. No overlap exists between the proteins phosphorylated through the two pathways (TSH and EGF/phorbol esters) (62).

The divergence of the two pathways, TSH/cAMP and that of the EGF/kinase C, also extends to newly synthesized proteins. Only one of the 26 proteins involved, proliferating cell nuclear antigen, is synthesized in response to TSH, EGF,

and phorbol esters. There are, however, differences in the kinetics of proliferating cell nuclear antigen synthesis—early through the TSH/cAMP pathway, and delayed through the other two pathways (63).

TSH induces rapid but short-lived increases in *c-myc* transcripts, whereas *c-fos* transcription is slightly delayed. EGF induces *c-myc* gene transcription later than does TSH. The kinetics of *c-fos* messenger RNA (mRNA) synthesis are, however, similar for TSH, EGF, and phorbol esters (65, 66). After an initial increase in *c-myc* mRNA in response to TSH, these transcripts decrease below control values, apparently related to a negative transcriptional signal (67). *c-myc* Gene activation is thus tightly regulated in its role in thyrocyte growth.

III. Epidemiological and Clinical Considerations

Thyroid carcinoma is an infrequent tumor with geographic variation in its prevalence. The annual rates of newly diagnosed cases vary from 0.9/100,000 men and 2.4/100,000 women in Britain (68) to 2.1/100,000 men and 5.2/100,000 women in the United States (69). Thyroid cancer accounts for 0.6% and 1.6% of all cancers among men and women, respectively, in the United States (70). In Japan the combined estimate is 1.4/100,000 of the population (71), whereas the crude relative frequency of thyroid cancer among Kuwaiti females was reported as 10.5% (72).

The incidence of thyroid cancer in the United States increased during the four decades 1935–1975, reflecting better diagnosis as well as the emergence of radiation-related cancers (73–75). That trend has stopped (76). Moreover, since the 1960s, survival rates for white patients have steadily increased (77).

Papillary carcinoma accounts for 85% of differentiated thyroid follicular cell cancers in iodide-sufficient countries (78), while the remaining epithelial thyroid cell tumors are predominantly follicular carcinomas. Anaplastic carcinoma, whose incidence has been decreasing, often arises from preexisting well differentiated (usually follicular) carcinoma (79, 80). An increase in the prevalence of follicular carcinoma under conditions of iodide deficiency or endemic goiter accounts for the overall increased incidence of thyroid cancer, including anaplastic carcinoma, in such areas (81–84a). Incidental microscopic (occult) to small papillary carcinomas are detected with such high frequency at autopsy (85) that one must conclude that the vast majority are not clinically relevant. The factors determining the transition from microscopic foci to detectable thyroid carcinoma are unknown.

Both cohort and case-control studies have established a strong link between external radiation and benign as well as malignant thyroid tumors (78, 86–99). By contrast, internal radiation probably does not enhance the risk of thyroid cancer (88, 100–102).

The relationship of thyroid cancer to endemic goiter has long been suspected (103). Comparative studies of thyroid cancer incidence between areas with and without endemic goiter have not, however, always supported that notion (104, 105). The effect of iodide supplementation on thyroid cancer incidence has also yielded conflicting results (106). Indeed, a recent case-control study from Italy suggested that the influ-

ence of iodide supplementation on cancer incidence was marginal (105). It is unclear whether the association of thyroid cancer with benign nodules and goiter is real or related to ascertainment bias (89, 92, 94). Likewise, the notion that thyroid carcinoma is more frequent in patients with Graves' disease, and tends to be more aggressive (107–109), has been challenged (110, 111).

Several host factors that determine the outcome of thyroid cancer have been identified. These include patient age, tumor size, histological appearance, local invasion, lymph node and distant metastases, and aneuploidy (112–114). Several clinical staging and prognostic systems have been based on these factors.

IV. Genetic Factors

Genetic factors are not generally thought to be important in predisposing to thyroid follicular carcinoma. Thyroid carcinoma is, however, increased in such syndromes as Gardner's syndrome (adenomatous polyposis) (115–121) and Cowden's disease (hamartomas) (122). In addition, reports of remarkable aggregation of papillary thyroid cancers in families (123–125) indicate that genetic factors may be relevant to thyroid follicular cell cancer. In a genetically homogeneous population, 3.8% of patients with papillary cancer had similarly affected family members (126). Genetic factors appear to be more important in the genesis of papillary carcinoma (123, 124, 126) compared to follicular carcinomas (see Ref. 127 for review; 128). Familial aggregation of follicular carcinoma is documented in families with dysmorphogenesis (129). Female patients with papillary carcinoma (but not their relatives) are at greater risk for other cancers [including breast, renal, and central nervous system malignancies (126, 130–133)].

There are also indications that certain ethnic groups may be more susceptible to the effect of ionizing radiation. Furthermore, in instances where siblings were irradiated, the incidence of thyroid tumor development was more than could be accounted for by chance (134).

The association of thyroid cancer with certain HLA alleles represents an interplay between the environment and genes in its pathogenesis, and has been found only in some populations. While no association of HLA antigens with well differentiated thyroid cancer was found in patients from an iodide-sufficient area (Newfoundland) (135), HLA-DR1 was significantly increased in patients from iodide-deficient Eastern Hungary (136, 137). The increase in HLA-DR1 was greatest in patients with follicular cancer (137). Sridama and colleagues (138) in Chicago reported an association of thyroid cancer and HLA DR7, particularly in patients with follicular carcinoma (138). Moreover, immunoglobulin G heavy chain allotypes interacted with HLA to enhance the risk conferred by the latter (139).

To account for these variations in HLA association with thyroid cancer in different geographic locations, it has been suggested that HLA DR1 positive individuals were at increased risk for thyroid cancer in iodide-deficient areas and that this predisposition disappeared once iodide sufficiency predominated (127, 137). In the absence of iodide deficiency,

other HLA-linked influences were important for those exposed to other environmental hazards. It is interesting that the susceptibility of rats to diethylnitrosamine carcinogenesis is related to the inactivation of recessive genes that map to MHC (140) and that nitrosamines induce thyroid tumors in mice (see also Section VIII.A.).

Last, the 3-fold greater incidence of thyroid cancer in females is probably related to the female hormonal milieu rather than to the contributions of active genes on the X chromosome. A case-control study (74) has suggested an association of thyroid cancer with pregnancy.

The cellular genetic events leading to thyroid cancer are somatic in nature and thus restricted to thyroid tumor tissue. The influence of paternal imprinting (141, 142) on oncogene activation is unknown.

V. The Cell Cycle and Cancer

Multicellular organisms have highly coordinated mechanisms to control cellular interactions. These signaling networks mediate normal embryonic development and the response to wound healing or infection. Aberrations in growth factor signaling pathways are intimately linked to cancer (143). Malignant cells arise as a result of a stepwise progression of genetic events that include the unregulated expression of growth factors or components of their signaling pathways.

Individual cells receive two classes of signals. One type promotes cell growth largely through the elaboration of growth factors. Another type of signal allows cells to inhibit the growth of their neighbors through regulatory proteins. The known number of genes involved in control of the cell cycle is increasing rapidly. When the function of some genes are lost, cells become unresponsive to growth-inhibitory signals. Several of these genes meet the requirements for "tumor suppressors." The inhibitory signals may involve not only proteins that function as switches in the cycle, but also diffusible growth inhibitors and hormones (144-146). The cell responds to inhibitory signals in three ways: 1) pausing in a certain phase of exponential growth often representing arrest at the G₁-S phase transition; 2) postmitotic differentiation; 3) senescence or apoptosis (programmed cell death) (147).

Growth signals allow cells in the resting (G₀) phase to enter and proceed through the cell cycle. The resting cell is first advanced into the G₁ phase of the cell cycle by "competence" factors, traverses the G₁-S phase and becomes committed to DNA synthesis under the influence of "progression" factors. Although the notion of "competence" was only demonstrated in some systems, its generalization is probably justified. Transition through the G₀-G₁ phase requires sustained growth factor stimulation over several hours. Disruption of the signal for only a short period of time results in the cell reverting to the G₀ phase. There is also a restriction point in the G₁ phase during which the presence of both competence and progression factors are necessary for the cell to advance through its cycle (Fig. 1). Thereafter, only progression factors are needed. Competence factors include EGF and FGF whereas IGF-1 and insulin act as progression factors (143). The cytokines such as TGF β , interferon- γ , and tumor




FIG. 1. The cell cycle. Sites of action of protooncogenes and growth and growth-inhibiting factors. Some protooncogenes, e.g. *c-ras* and *c-erbB*, contribute to the cell "competence" and are complemented by the action of others, e.g. insulin, to allow progression of the cycle beyond the G₁-S restriction phase, thus committing the cell to DNA synthesis. For human thyroid cells TSH is a "competence" factor. Both retinoblastoma (Rb) and p53 proteins act at the G₁-S restriction phase to halt cell cycle progression. One of the pathways of Rb growth-inhibiting influence is through the induction of the TGF β gene [Adapted with permission from S. A. Aaronson: *Science* 254:1146-1153, 1991 (143). ©1991 by the AAAS.]

necrosis factor can antagonize the effects of growth factors. Activated oncogenes may encode growth factors (e.g. *c-sis*), receptor protein kinases, or other enzymes that participate in mitogenic signaling. Oncogenes such as *c-ras* and *c-erbB* function as competence factors. Other oncogenes, e.g., *bcl-2* (148-150), act by blocking apoptosis and allowing cells to proliferate preferentially in response to mitogenic signals and may indeed block the apoptotic effects of *c-myc* (151-153). p53 And retinoblastoma (Rb) proteins function by regulating the cycle at the critical G₁ to S phase transition (154-156). Several of the growth factors participate in tumor progression by providing in a paracrine fashion the environment for proliferation, invasion, and metastases of epithelial cells. A number of growth factors including FGF, EGF, and hepatocyte growth factor participate in tumor-sustaining neoangiogenesis.

Are tumor cells intrinsically more unstable than normal cells, or do they merely acquire their abnormal karyotypes through increased division and intense selection at normal rates of chromosomal rearrangement? This critical question has been definitively answered only recently (157). Many tumor cells exhibit rates of gene amplification several orders of magnitude greater than normal cells. Gross chromosomal changes such as marked aneuploidy, translocations, or dele-

tions are not nearly matched by the rates of point mutations (158). Such genomic instability is a genetic trait inherited in human cancer-prone syndromes (159, 160). These may be associated with a defect in G₁-S phase and is upstream from the p53 gene in a pathway that activates the G₁-S checkpoint. X-ray exposure of eukaryotic cells results in G₁ arrest and is associated with an increase in p53 protein. Cells lacking p53 or having a dominantly negative mutation lack this G₁-S delay (161). In premalignant cells, increased genetic instability after DNA damage related to radiation, carcinogens, or single strand breaks results in duplication, amplification, and replication of broken sister chromatids (157, 161). Some of these events may result in the loss of growth suppressor genes and amplification of oncogenes, although the picture is likely more complex (162, 163). This route for carcinogenesis is worthy of study in radiation-related thyroid tumors.

VI. Thyroid Cell Cancer—Mono or Polyclonal?

Recent work involving mutated *ras*, *GSα* (164), and rearranged RET/PTC (PTC = papillary thyroid carcinoma oncogenes) (165) have indicated variations in the presence of mutant gene products within the same tumor tissue. Whether these abnormalities are secondary or primary in origin, or related to multiple foci each arising with a separate array of mutations, are pertinent questions, and raise the issue of the clonality of thyroid tumors. If tumors arise by a series of rare mutations or epigenetic events, it is implied that such lesions derive from a single cell.

Determination of clonality can be achieved by X chromosome inactivation analysis, taking advantage of the fact that one of two X chromosomes is functionally inactivated early in mammalian embryogenesis (166). X-inactivation is random involving either the paternal or maternal X chromosome so that female tissues are mosaics containing an equal mixture of cells in which maternal or paternal chromosomes have, respectively, been activated. Taking advantage of differences in restriction fragment length polymorphisms (RFLPs) between two alleles in heterozygote females and the ability of some restriction enzymes to identify the methylation differences (167) between the functional and inactivated genes, the clonality of benign and malignant thyroid tumors has been studied (168–171). Using RFLPs at the hypoxanthine phosphoribosyltransferase, phosphoglycerokinase (PGK), and more recently M27 marker genes (171), most benign nodules and malignant thyroid tumors were shown to be monoclonal in origin. Nine of 13 rapidly growing nodules within 12 multinodular goiters were also found to be monoclonal using the highly informative M27 marker (171). One hyperfunctioning thyroid nodule in a 22-month-old child was polyclonal (172).

A problem in this analysis of solid tumors is the potential contamination of tumor samples by nontumorous tissue. The polymerase chain reaction (PCR) may facilitate the analysis for monoclonality of small homogenous tissue fragments (173): *HpaII* digestion of DNA followed by PCR amplification with primers on each side of *HpaII* restriction site within the PGK gene would allow the amplification of only the methylated copy of the gene (174). Random inactivation of PGK

is associated with two DNA fragments whereas monoclonality is associated with one band.

VII. Growth Factors and Their Receptors in Thyroid Tumors

Unregulated growth signals acting on cells with multiple transforming lesions result in malignancy. These signals may be the result of constitutive synthesis of growth factors, constitutive activation of their receptors, or of switches in the signal transduction pathways. It is thus of interest to review what is known in this context of thyroid carcinoma (175).

EGF and EGF receptor have been detected by immunohistochemistry in malignant thyroid tumors, but not in normal thyroid tissue, benign tumors, or multinodular goiter (176, 177). The presence of both the growth factor and receptor was more frequent in papillary than in follicular carcinoma samples (176). EGF receptor was also detected by radioligand assays in normal and hyperplastic thyroid tissue, as well as benign and malignant tumors (178). In that study, the level of EGF receptor was found to be greatest in anaplastic tumor tissue (178). The predominance of EGF receptor expression in papillary carcinoma was extended to the transcripts of *c-erbB1* and *c-erbB2/neu* oncogenes, which encode EGF receptor or analog, respectively (179, 180). A 2- to 3-fold increase in *c-erbB1* and *c-erbB2/neu* mRNA in papillary carcinomas and their lymph node metastases, as well as in one benign adenoma, compared to thyroid tissue were reported (180). No structural abnormalities in these protooncogenes were detected in a large number of benign and malignant thyroid tumors (179, 181).

EGF expression was shown to be of prognostic value in that tumors with higher EGF expression were more likely to recur (177), analogous to the findings with *c-erbB1* and *c-erbB2/neu* (180).

No evidence for rearrangement or mutation of *c-erbA* (T₃ receptor) was found in thyroid tumors (179). At least some mutated T₃ receptors interfere with T₃ action (as well as perhaps that of related receptors in a dominant manner (Refs. 182–184 for review), which makes them potential candidates for tumor suppressors (147).

Analysis by immunohistochemistry revealed TGFβ expression in 58% of malignant thyroid tumors, but not in benign adenomas or normal thyroid epithelium (185). In follicular carcinomas TGFβ immunostaining was related to the presence of H-*ras* 61 (Glu → Arg) mutation. The fact that TGFβ is a potent inhibitor of epithelial thyroid cell growth and is synthesized by thyroid cells suggests that the increase in TGFβ may compensate for decreased TGFβ sensitivity (186, 45). TGFα appears to regulate the expression of EGF receptor in some tumors and thus promotes their growth (187), but was not studied in thyroid cancers. Apparently both benign and malignant thyroid tumors elaborate IGF-I (38, 188) more than normal tissue, which may allow nodules to become autonomous (26).

In anaplastic thyroid carcinoma, receptors not normally expressed in the normal thyroid gland may be detected. Functional but aberrantly glycosylated receptors for platelet

derived growth factor (PDGF) were described in an anaplastic thyroid carcinoma cell line (189).

VIII. Oncogenes Involved in Thyroid Carcinogenesis

A. *ras*

The mechanisms by which receptor tyrosine kinases stimulate specific intracellular signaling pathways have been elucidated by the identification of a conserved protein motif of the *src* homology 2 (SH2) domain, which is found in a remarkably diverse group of signaling proteins (190). This includes nonreceptor protein kinases, phospholipase C- γ 1, *ras* guanine triphosphatase (GTPase) activating protein (GAP), and other molecules referred to as adaptors. Proteins with SH2 domain frequently have another distinct sequence, the SH3 domain, which is also implicated in the regulation of protein-protein interaction during signal transduction. For example, the autophosphorylation of the insulin receptor results in tyrosine phosphorylation at the SH2 consensus sequence of a 185 kDa cytosolic protein (27), which then recruits and activates SH2-containing phosphatidylinositol (PI) 3'-kinase (191). A similar scheme may be envisaged for GAP, which upon phosphorylation associates through SH3 domains with guanine nucleotide exchange factors (190, 192), thus regulating *ras* and *ras*-like G proteins. Moreover, tyrosine protein kinases, through SH2-containing protein adaptors (193, 194) may activate *ras*-dependent signaling pathways (Fig. 2). The intermediate elements in this cascade have been recently elucidated, at least in some settings (see Ref. 195 for review).

GAP stimulates the GTPase activity of *ras* which is a critical component of intracellular mitogenic signaling pathways (194, 196) and also acts as a negative regulator of *ras* function (197). Mutations that cause oncogenic activation of *ras* lead to accumulation of *ras* bound to GTP, the active




FIG. 2. A postulated schema for the role of *src*-homology (SH) domains 2 and 3 for coupling tyrosine kinases to G proteins. According to this model, proteins with SH2 and SH3 consensus sequences and catalytic domains would be capable of multiple point interactions upstream with tyrosine kinases and downstream with *ras*-like small G proteins (adaptors) and their substrates. Interaction with SH3 domains allows GTPase activating proteins (GAPs) to recruit guanine nucleotide exchange factors regulating *ras* and adaptor proteins (190).

form of the molecule (196). These mutations block the ability of GAP to promote conversion of *ras* to its inactive, GDP-bound form (194). GAP may also function in a complex with *ras* as an effector of its downstream signaling functions (198). Thus, mutations that impair interaction of *ras* with GAP also block the biological function of *ras*. *ras* Protein (p21) is probably excited by an activator upstream in its signaling pathway and passes these signals to a downstream effector pathway (Fig. 3). It is conceivable that constitutive activation or amplification of the signals acting along *ras* or *ras*-like pathways may be involved in cell transformation. It is also possible that mutations in the SH2 adaptors may have similar consequences. The signals involved in the increased expression of *ras* during normal thyroid cell growth (199) are unknown.

ras Protooncogene mutations are found in more than 30% of human tumors (200). Three families of *ras* proteins have been identified, Ha-*ras*1, K-*ras*2, and N-*ras*, each of which is located in a separate chromosomal region (196). All three are subject to mutations, although some mutations tend to predominate in specific tumors, e.g. K-*ras* mutation at codon 12 in pancreatic carcinoma (201) and adenocarcinoma of the lung (202). Mutations at residues 12, 13, or 61 of any of the *ras* protooncogenes convert them into active oncogenes (196); other mutations may also be relevant to constitutive signaling. Mutations at codon 61 are the most efficient in changing




FIG. 3. The role of *ras* in the signaling pathway from tyrosine kinases and other receptors to transcription factors. Raf-1 autokinase is an important station in the *ras*-mediated transduction of the action of many tyrosine kinases—including *src*. It is apparent, however, that the maximal activation of Raf-1 is dependent on synergy between a direct as well as a *ras*-mediated effect. Mitogen-activated protein (MAP) kinases are positioned in the *ras*-pathway leading to the phosphorylation of the ribosomal S6 kinase (pp90^{rak}) and in turn nuclear transcription factors such as *c-fos* and *jun*. MAPK is activated by a cascade of MAPkinase kinase (MAPKK) and MAPkinase kinase kinase (MAPKKK). One of the targets of phosphorylation by EGF treatment of thyroid cells is MAPK (56). The details of this activation pathway are, however, unclear. Platelet dependent growth factor (PDGF), which is expressed aberrantly in anaplastic carcinoma (189), activates the *ras* and Raf-1 pathways (143). Cellular elements intermediary between receptor tyrosine kinase—Sem2 and son of sevenless (*sos2*) which interact through SH2 and SH3 domains—and *ras* have recently been identified (195). [Adapted with permission from T. M. Roberts: *Nature* 360:534–535, 1992 (225). © 1992 Macmillan Magazines, Ltd.]

the conformation of *ras* proteins and are associated with a reduction in GTPase activity (203).

ras Mutations are not sufficient for malignant transformation. The ability of v-Kris-*ras* to transform rat (204) and Ha-*ras* human (205) thyroid cells in primary cultures seemingly contradicts the role of *ras* mutations as early events in thyroid carcinogenesis. Helper viral elements, however, do complement the role of *ras* in transformation in these studies (13). Indeed, even rare events such as double *ras* mutations are not necessarily enough for thyroid cell transformation (206). Moreover, a mutant *ras* gene expressed from a retroviral vector induces thyroid cell proliferation only in the presence of growth factors (207). This growth was inhibited by phorbol esters through regulation of protein kinase C activity (207). Overexpression of the *ras*-21 protooncogene appears to be part of growth-promoting pathways in normal human thyroid and nodular goiter (199).

Distribution of *ras* RFLP in thyroid cancer DNA specimens showed unusual patterns, some of which probably reflected mutation in the coding sequence of the gene (208, 209). An increased abundance of *ras* protein or mRNA was reported in some studies but could not always be related to the degree of tumor de-differentiation (176, 210, 211). Rearrangement of *ras* protooncogenes in thyroid tumors is uncommon and not specific for malignant tumors (179).

ras Gene point mutations are of more interest: mutations in all three families of *ras* oncogenes have been detected in both benign and malignant thyroid tumors (164, 179, 206, 212-223). Most of the mutations were found at residue 61 of H-*ras* and N-*ras* genes (206, 212, 220). Variation in the prevalence of *ras* point mutation thyroid tumors does occur among different series (164, 206, 220) and may be related to the histological and degree of differentiation of the tumors as well as genetic determinants and environmental factors, such as dietary iodide supply (206). It is of interest that thyroid tumors induced by chemical carcinogens in rats involve activation of H-*ras* (219) whereas Ki-*ras* is activated exclusively in 60% of radiation-related tumors in man (224) and rat (219). In the former, radiation does not appear to enhance the overall incidence of *ras* protooncogene mutations (224).

ras Mutations occur predominantly in follicular thyroid cancers (206, 212, 220, 223). Twelve percent of Hurthle cell tumors, a variant of follicular carcinoma, harbored N-*ras* mutations (215). Interestingly, normal tissue adjoining the tumor also exhibited *ras*-activating mutations, emphasizing that cellular events in addition to *ras* mutations are necessary for malignant transformation. The notion that *ras* mutation is relatively specific for microfollicular benign adenomas as opposed to macrofollicular histology (220) could not be sustained (206). An increased frequency of *ras* mutation in metastatic tumors was observed in some (179, 220) but not in other series (206, 214).

The *ras* mutations reported are generally of the transition type (in which a purine is substituted for a purine or a pyrimidine for a pyrimidine), although it has been suggested that transversion mutation (in which a purine is substituted for a pyrimidine and *vice versa*) may be relevant to dediffer-

entiated thyroid tumors (220). Preferential mutation at H- and N-*ras* 60 is apparently lost in anaplastic carcinoma (220) and awaits further confirmation.

B. *c-myc* and *c-fos*

The *c-myc* nuclear proteins act as transcription factors. *c-myc* Dimerizes with its partner *Max* protein and binds to core consensus DNA sequence (226-229). *Max* heterodimerizes preferentially with *myc*, which homodimerizes poorly (230). The formation of *myc/Max* dimers is essential for *myc* transforming activity whereas *myc* homodimers are inactive (231). *c-myc* Synthesis declines as the cell cycle progresses (232) and is shut off with inhibition of proliferation associated with differentiation. *c-myc* Has also a central role in some forms of apoptosis (233-235), particularly under growth-limiting conditions. *Bcl-2* mitigates the apoptotic effects of deregulated *myc* expression without affecting its ability to promote continuous cell growth (152, 153).

Alterations in the c-MYC locus occur in a variety of tumors primarily leading to constitutive expression of *myc*. Such deregulation may result in a shift from *Max* homodimers to *myc-Max* heterodimers, and enhances the activation of genes normally modulated during growth and differentiation. Both *myc* and *Max* may be regulated by phosphorylation (236).

Activation of *c-myc* by translocation is restricted to lymphoid tumors. *c-myc* Amplification appears to be a secondary selection factor for increased transformation, as opposed to a primary genetic lesion leading to malignancy (237). The 5'-deletions of the *c-myc* gene described in thyroid tumors (238, 239) were subsequently found to be germline in origin (238) and to exist in unaffected healthy individuals. No gross rearrangements of *c-myc* were documented (214, 240-242). Immunohistochemical studies and mRNA abundance indicated increased expression in thyroid tumor tissue (176, 177, 240, 242, 243). The abundance of *c-myc* transcripts may predict the course of thyroid malignancy: the less differentiated a tumor, the greater the abundance of *c-myc* transcripts (240, 244, 245). Since *c-myc* message abundance was not reflected (at least in papillary carcinoma) in the level of *c-myc* gene products (177), there may be differences in *c-myc* message stability in such tumors. *c-myc* mRNA correlated negatively with that of the TSH receptor (244, 245).

The protooncogenes *c-fos* and *c-jun* are immediate early genes that regulate the expression of specific target genes (246). Depending on cell type the expression of these genes can be activated by second messengers which stimulate protein kinase A and C activities and calcium flux (247-250). The protein products of the genes interact through leucine zipper domains to form *fos/jun* heterodimers or *jun/jun* homodimers. As components of the AP-1 transcription complex they bind to the regulatory sequence of many genes to initiate transcription (251).

No evidence for rearrangement or amplification of *c-fos* was noted in thyroid tumors (214, 242). Increased *c-fos* transcripts were found in 60% of malignant tumors and 90% of benign adenomas. There was no relationship between the increased transcripts to the biological behavior of the tumors (240). Similar conclusions were reached in our studies of 42

th of di
C.
or
ki
ur
ar
ar
pr
pe
ca
tra
de
tif
va
th
tra
ki
tic
te
re
m
th
(F
be
ar.
ki
(2
is
sp
C
de
86
Fr
in
pl
fr
wi
89

thyroid tumors (our unpublished observations). This pattern of *c-fos* message expression is consistent with its role in differentiation.

C. PTC/RET oncogene

PTC/RET [previously PTC for papillary thyroid carcinoma] oncogene (252-256) results from the fusion of the tyrosine kinase domain of the protooncogene *ret* with a functionally unknown sequence as a result of chromosomal rearrangement. The most frequent partner of this rearrangement is the D10S170 locus (257-259) identified with probe H4 (252). The activation of PTC/RET was first suspected by Fusco *et al.* (260) when they found 25% of papillary carcinoma samples were positive in a transfection assay. The transforming activity was cloned and identified by two independent groups (252, 254). The rearrangement was identified by RFLP and the breakpoint was found to be highly variable within the H₄ sequence but relatively constant in the *ret* protooncogene, mapping to the intron between the transmembrane domain and the first exon of the tyrosine kinase domain (252, 261). All the characterized recombinations, however, produce identical transcripts encoding a protein of 520 residues (252, 257). The activation of RET/PTC results from paracentric inversion of the long arm of chromosome 10, inv(10)(q 11.2 q21) with breakpoint involving the regions where RET and D10S170 are located (262) (Fig. 4).

The unrearranged *ret* protooncogene sequences appear to be involved in neuronal differentiation (263) and to be amplified in neuroendocrine tumors (264, 265). Tyrosine kinase is not activated in the unrearranged *ret* gene product (266), although its natural ligand is unknown. There are two isoforms of *ret* protooncogene which arise by alternative splicing and differ by having 9aa vs. 51aa at their respective C termini. The 51aa variant is glycosylated to a variable degree in different cell types (267-270). The *ret* protooncogene is not expressed in thyroid follicular cells but is in

parafollicular cells, while its rearranged form is expressed in papillary cancer cells (252).

The RET/PTC rearrangement is specific for the thyroid, being undetectable in 250 nonthyroidal malignancies (165, 261, 271). Rearrangement of *ret* previously described in other tumor DNA was found to have occurred *in vitro* during transfection procedures (256). There is a wide geographic variation in the rate of RET/PTC rearrangement in thyroid tumors (Table 1) perhaps related to genetic and/or environmental factors as well as methodology (165, 261, 271, 272). Transfection assays (252, 260), RFLP (252), and reverse transcription-PCR (RT-PCR) (165, 261, 271, 272) have been used to examine oncogene activation and rearrangement. Only the RT-PCR approach is predicated on the identity of the sequence bridging the breakpoint. The rearrangement is apparently specific for papillary carcinoma, although it is unlikely to occur in more than 20% of these tumors (271, 272).

Two groups have, however, reported RET/PTC rearrangement in benign nodules, nodular goiter, and follicular carcinoma (165, 274). In the study of Ishizaka *et al.* (165), protooncogene activation was regionally localized within benign nodules and multinodular goiter and was attributed to the high rate of microscopic papillary carcinoma in Japan. The issue of focal PTC/RET rearrangement in such microfoci may be resolved by *in situ* hybridization with an appropriate probe (275). In a subsequent report, Jhiang *et al.* (261) do not comment on the frequency of PTC/RET rearrangement in follicular adenomas they reported earlier (274). It is interesting that *ret* is rearranged in tumors with follicular/papillary histology (275), giving molecular credence to the designation of these tumors as papillary.

It has been suggested (261) that patients with papillary carcinoma with rearrangement of the *ret* protooncogene have a greater likelihood of developing distant metastases. These studies in a small group await confirmation.

It also bears emphasis that only one third of the samples with *ret* rearrangement show evidence of fusion with H₄ (271). The remaining samples will clearly *not* be identified by an RT-PCR strategy in which a probe spanning the breakpoint is used for screening (261, 271, 272).

The issues of specificity, the mechanisms of RET/PTC activation, and the rearrangement partners other than H₄ are being rapidly clarified (266). *ret* May undergo rearrangements in papillary thyroid carcinoma with a partner other than D10S170 (H4) sequences. This rearrangement may involve either the 9aa (Ptc2) or 51aa (Ptc1) C-terminal form of *ret*. All forms of *ret* rearrangement are distal to the transmembrane domains and therefore localize the hybrid protein intracellularly (Fig. 5). As a consequence, the rearranged gene product constitutively activates the tyrosine kinase, which autophosphorylates at a tyrosine residue. The ptc1 and ptc2 are apparently regulated by phosphotyrosine phosphatases (266). The translocation of the activated rearranged *ret* oncogene may allow them to escape regulatory influences and to constitutively signal mitogenic pathways. It remains to be seen what proportion of papillary carcinoma utilizes non-D10S170 sequences as a partner in the PTC/




FIG. 4. Paracentric inversion of the long arm of chromosome 10 (10 inv(10)(q11.2 q21)) in thyroid papillary carcinoma. This inversion places an unknown sequence detected by probe D10S170 upstream from the tyrosine kinase domain of the *ret* protooncogene. [Reproduced with permission from M. A. Pierotti *et al.*: *Proc Natl Acad Sci USA* 89:1616-1620, 1992 (262).]

TABLE 1. Geographic distribution of PTC rearrangement

	USA ^a	USA ^b	France ^a	Italy ^a	Japan ^c	Saudi Arabia ^d
Papillary	11/65	4/32	8/70	14/42	1/11	1/40
Follicular	0/11	0/3	0/13	0/13		0/4
Anaplastic	0/2		0/5	0/8		0/5
Adenoma		0/8	0/18	0/16	4/19	0/1
Others		0/28			1/2	0/7


^a Santoro *et al.* (271).^b Jhiang *et al.* (261).^c Ishizaka *et al.* (165).^d Zou *et al.* (272).

FIG. 5. The rearrangements of the *ret* protooncogene in papillary thyroid carcinoma. The genomic *ret* gene is shown schematically at the top of the figure. Two forms of the *ret* protooncogene arise by alternative splicing and have in their COOH termini nine or 51 residues, respectively. The 51 COOH terminus form is glycosylated to variable degrees and the degree of glycosylation may differ from one tissue to another. In the thyroid, 150-kDa and 170-kDa differentially glycosylated forms exist. In thyroid papillary carcinoma, both *ret* protooncogene isoforms are rearranged. Initially, the 51aa. COOH terminus form was found rearranged fusing its kinase domain with an unknown sequence H4 (D10S170) (Ptc1); this probably accounts for no more than 30% of the rearrangement. Other sequences, yet to be identified, can also be involved in the *ret* protooncogene rearrangement and result in an 81 kDa protein (ptc2). Apparently, the short isoform of the *ret* protooncogene undergoes a similar type of rearrangement in papillary carcinoma (shown here as ptcx, a 76 kDa hybrid). All rearranged forms of *ret* protooncogene lose their transmembrane (TM) domain and the fusion proteins are thus localized intracellularly. Whether because they undergo conformational changes or are exposed to new substrates, these fusion products exhibit constitutive activation of the *ret* tyrosine kinase. (◻ = carbohydrate adducts).

RET rearrangement (266, 271). RET/PTC oncogene-transfected rat thyroid cells lose their differentiated function but are unable to grow in soft agar or to cause tumors in nude mice. An undifferentiated malignant phenotype is obtained when both RET/PTC and *Ha-ras* or *Ki-ras* oncogenes are transfected (276).

We have studied a large number of benign and malignant tumors using the RT-PCR approach (272). Only one out of 40 thyroid papillary carcinomas was found to have undergone PTC/RET rearrangement, suggesting a very low rate of rearrangement RET/D10S 170 locus in the Saudi population (Table 1). Of interest was that this tumor also harbored a mutation in the tumor suppressor gene p53 (Ala¹⁶¹ → Thr). It is unclear whether this is a chance finding or implies selection of *ret* rearrangement for p53 mutation.

Given the activation of *ret* protooncogene by rearrangement in papillary thyroid carcinoma, it is particularly

intriguing that missense germ-line mutations of RET have recently been proposed as the genetic basis for MEN2A liability (273).

D. TRK-T1

A second transforming event involving gene rearrangement which activates a tyrosine kinase protooncogene was found in papillary carcinoma (277). The oncogene involved, called TRK-T1, is generated by an intrachromosomal rearrangement that links the tyrosine kinase domain of TRK protooncogene to the 5'-region of the TPR gene, both mapping to chromosome 1 q23-q24 (278, 279). The TRK protooncogene encodes a surface receptor for nerve growth factor (NGF), and the TPR gene a protein with likely cytoskeletal function. The mechanism of TRK-T1 activation is not well understood and may include TRK rearrangement

that
the c
in cc
inter
anisi
M
recei
nom
be c
natic
tivat

E. M

Ti

two

is he

an e

men

whc

mat

lula

The

ter

tran

cytc

in u

(rev

H

of a

whc

of j

rece

the

regu

kina

T

cati

var

sub

es's

agg

sue

in z

care

neu

A

illat

off

or t

disc

cati

arra

δ-s

am

det

cell

the

feri

that activates the gene, moves the hybrid gene product to the cytoplasm exposing it to unusual substrates or may result in conformational changes which mimic those produced by interaction with the ligand, or combinations of these mechanisms (280).

More than one type of rearrangement involving the NGF receptor chain may be generated in papillary thyroid carcinomas (280). The TRK-T1 rearrangement was suggested to be common in papillary carcinoma (277), which, in combination with *ret* rearrangement, accounts for constitutive activation of tyrosine kinase in 50% of these tumors (280).

E. Met

The *met* oncogene is a 190-kDa heterodimer composed of two disulfide-linked (α , β) subunits. The α -subunit (50 kDa) is heavily glycosylated and extracellular. The β -subunit has an extracellular domain involved in ligand binding, a transmembrane segment and cytoplasmic tyrosine kinase domain whose activity is regulated by phosphorylation (281). The mature receptor is derived from glycosylation and extracellular cleavage of a 170-kDa precursor by urokinase (282). The multifunctional cytokine hepatocyte growth factor/scatter factor (HGF/SF) is the ligand for *met*. Multiple *met* transcripts that encode proteins differing in extracellular and cytoplasmic domains have been identified: one form results in uncleaved precursor chain and another in a soluble form (reviewed in Refs. 281 and 283).

HGF/SF is itself a heterodimer arising from the cleavage of a precursor. The light chain is related to serine protease whereas the heavy chain belongs to the plasminogen family of proteins (284-286). HGF/SF binding to *met* enhances receptor kinase activity by tyrosine autophosphorylation of the β -subunits whereas receptor kinase activity is negatively regulated by serine phosphorylation mediated by protein kinase C.

The *met* oncogene is constitutively activated by amplification of the gene or through the expression of a splice variant in which the precursor is not cleaved into the two subunits (281). Activation of *met* is associated with mitogenesis as well as motogenesis and may thus contribute to tumor-aggressive and metastatic behavior in neoplastic human tissues (285, 286). *Met* transcripts and protein were increased in a high proportion of gastrointestinal and hepatocellular carcinomas, and in malignancies arising from meningeal and neural tissues.

Met oncogene is amplified in approximately 70% of papillary and poorly differentiated carcinomas, but in only 25% of follicular carcinomas. *Met* was not detected in anaplastic or medullary carcinomas, nor in a variety of benign thyroid disorders and normal thyroid tissue (287, 288). The amplification of *met* gene was not associated with gene rearrangement, although several samples exhibited an 85-kDa β -subunit soluble variant (288). In most tumors with *met* amplification an autocrine source of HGF/SF is not readily detectable. The elaboration of this ligand by parafollicular cells (289) seemingly establishes a paracrine relationship in the thyroid. Interestingly, of the 10 papillary or poorly differentiated carcinomas also examined for RET and TRK re-

arrangements, in three RET was rearranged and TRK in one and, except for one tumor with RET rearrangement, all samples exhibited amplification of *met*. This emphasizes the important contribution of *met* in the tyrosine kinase "pathway" for papillary carcinogenesis (288). *Met* overexpression was associated with aggressive clinical and histological phenotype.

Activation of tyrosine kinase, whether by gene amplification or rearrangement, appears to be highly specific for the transformation of thyroid follicular cells into papillary tumors. It is also of interest that papillary thyroid carcinoma is the only type of nonhematopoietic tumor with a high frequency of gene rearrangement.

F. Gsa mutations

The G proteins are a subfamily of the GTP-binding proteins, which include *ras* and *ras*-like proteins (290, 291). The G proteins are heterodimeric, composed of α -, β -, and γ -subunits each encoded by a distinct gene. The α -subunit shows structural and functional homology with other members of the GTP binding protein superfamily. It binds guanine nucleotides with high affinity and specificity and has intrinsic GTPase activity. β - and γ -subunits are noncovalently bound into a dimeric complex. They are necessary for regulating the function of certain α -subunits as well as in directing the trimolecular complex to the plasma membrane (reviewed in Refs. 292 and 293).

G proteins couple a diversity of receptors with their effectors by acting as molecular switches activated and deactivated by the GTPase cycle (292). The variety of G subunits identified, details of the functional domains of $G\alpha$, and an overview of the strategies used to elucidate the receptor/G protein coupling were recently reviewed by Spiegel *et al.* (292). $G\alpha$ is utilized widely as a positive transducer for the activation of adenylate cyclase and calcium channels. Cholera toxin ribosylates Arg²⁰¹ and thus impairs the hydrolysis of the γ -phosphate of GTP. Natural mutations at this residue (*gsp*) have an identical effect. Mutations at residues 232, 233, or 234 (numbering based on 394 residue form of $G\alpha$) have been shown either to inhibit activation of $G\alpha$ or its GTPase activity. The glutamine at position 227 corresponds to position 61 in *ras* p21.

Activating germline mutations of $G\alpha$ may well be lethal. The activating mutations described in McCune-Albright syndrome (294, 295) show tissue mosaicism, suggesting that they have arisen after the first rounds of fertilized ovum division. The $G\alpha$ -inactivating mutations noted in Albright's hereditary osteodystrophy may be less deleterious in early development. It is of interest that the activating mutations in McCune-Albright syndrome have been restricted to Arg²⁰¹, whereas multiple distinct (and heritable) heterozygous inactivating mutations were uncovered in families with Albright's hereditary osteodystrophy (292, 296, 297).

A subset of GH-producing pituitary tumors with high basal adenylate cyclase and which tend to be small (298) have been found to harbor α -subunit (*gsp*) mutations (299-301). The mutations involved predominantly Arg²⁰¹ \rightarrow Cys or Gln²²⁷ \rightarrow Arg or Leu. Similar mutations were described in a

small proportion of thyroid tumors (164, 300, 302, 303). Twenty five percent of follicular adenomas were found to harbor *gsp* mutations (298, 302, 303). There may be a predilection for these mutations in microfollicular adenomas. The rate of these mutations was found in a recent study to be less than 3% (304). Some of the mutations were novel, e.g. Gln²²⁷ → His. Mutations reported in papillary and follicular carcinoma are uncommon (300, 302, 304); the substitutions were also unusual (Arg²⁰¹ → ser, Gln²²⁷ → Lys and Gln²²⁷ → His) (302). As anticipated, *gsp* mutations were more common in tumors selected with high basal adenylate cyclase activity (302).

A recent study (164) is worthy of separate consideration, since it shows geographic variation (Germany vs. the United States) in the rates of *gsp* and *ras* oncogene activation in 32 differentiated carcinomas and regional differences in the activation of these two oncogenes in fragments of the same tumors. The authors suggest that in samples from iodide-deficient Germany *gsp* mutations are more common than *ras* (in direct contrast to US samples) and that these mutations are much more frequently identified in tumor section than from whole tumors. Because thyroid tumors are monoclonal (see above) some of these mutations are probably late events. They may, nevertheless, modify tumor biological behavior (164). The potential interaction between *ras* and *gsp* mutation in thyroid tumor requires further clarification.

To what extent regional variation within the same tumor accounts for variation in *gsp* mutation within iodide-sufficient regions is unclear (298, 302-304) and, indeed, given the number of samples studied, the differences between some of the studies are not statistically significant. Our own experience with a detailed study of 32 thyroid tumors uncovered no mutation at Arg²⁰¹ or Gln²²⁷. We have, however, found a mutation in exon 1 of *Gsa* (Lys³⁴ → Arg) in one follicular adenoma. The mutation is placed in a functional attenuator domain in *Gsa*. These experiments were predicated on the notion that N-terminal domains of *Gsa* regulate the rate of α s activation by guanine nucleotides independent of GTPase activity intrinsic to *Gsa*. It is likely a modulator domain of GDP dissociation and GTP activation (305). Whether this is a random somatic mutation or a normal polymorphism, and whether it alters *Gsa* function and is thus relevant to thyroid tumorigenesis, remains to be determined.

The potential oncogenic role of constitutive elevation of cAMP is emphasized by transfection of A₂ adenosine receptor cDNA into cells or its targeting for expression in the thyroid of transgenic mice under the influence of the thyroglobulin promoter (306, 307). In the latter case, constant adenylate cyclase activation is associated with thyroidal hyperplasia and nodular goiter (307). On the other hand, Gi_s2, a Gi_s subtype regulated by TSH (308), was found to be constitutively increased in autonomous adenomas, but not in thyroid cancers. The increase in Gi_s2 was associated with decreased cAMP levels in tissues tested (308).

IX. The TSH Receptor

The TSH receptor is a member of the seven-transmembrane segment G protein-associated receptors. In common

with the LH/CG and FSH receptors, the TSH receptor has a large extracellular domain with multiple glycosylation sites (see Ref. 309 for review). Structure-function analysis of the receptor has delineated some of the receptor functional domains (310, 311). The second cytoplasmic loop, COOH-terminal domain of the third cytoplasmic part, and amino terminal of the intracellular tail appear to be important in transducing the signal initiated by agonist binding (309, 310). The intracellular tail does not possess a consensus sequence for protein kinase A substrate but does have several potential kinase C substrate sequences (312). The TSH receptor third intracellular loop has a distinctive motif found in nonreceptor tyrosine kinases (312). Mutation of the tyrosine in this motif did not, however, alter receptor activity (310). Three major transcripts (4.5, 1.7, 1.3 Kb) were observed, probably resulting from alternate splicing of the mRNA. The two smaller variants can potentially result in soluble TSH receptor forms (313, 314). Two insertions (8aa at the NH₂ and 50aa at the COOH-end of the extracellular domain), are characteristic of the TSH receptor (see Ref. 309). The former is essential for TSH binding and receptor activation, whereas the biological relevance of the latter is unclear.

It is now possible to verify, at the molecular level, earlier observations of negative correlation between the degree of differentiation of thyroid tumors and their ability to bind [¹²⁵I]TSH and to stimulate adenylate cyclase (reviewed in Ref. 315). Benign thyroid tumors have a similar number of TSH receptor transcripts to those found in normal thyroid (244, 245). With decreasing degrees of differentiation, fewer TSH receptor transcripts are found. Indeed, a tolerable degree of negative correlation between the clinical staging of the disease and receptor mRNA abundance exists (244, 245), with some exceptions (244). With an increase in dedifferentiation, however, expression of thyroperoxidase is lost (245, 316) well before thyroglobulin transcripts (245). The loss of TSH receptor gene expression appears to be a late event in thyroid tumor dedifferentiation (245). We did not find a preferential expression of receptor variants in thyroid tumors (244). The relevance of a report of variation in the size of TSH receptor gene determined by RFLP in follicular adenomas (317) is unclear.

Increasing dedifferentiation of thyroid tumors and decrement in TSH receptor mRNA was associated with an increase in β -adrenergic receptor transcript numbers, deduced by quantitative PCR (318). It is unclear whether these mRNAs are translated into functional β -adrenergic receptors or whether they have undergone mutations that will render them constitutively activated. This area of investigation is worthy of further exploration, as site-specific mutations of the C-terminal part of the third intracellular domain of the α_{1B} -adrenergic receptor was associated with constitutive activation of PI hydrolysis (319) and neoplastic transformation. These mutants have not, however, been observed naturally.

That spontaneous mutations in the third intracellular loop of the TSH receptor are relevant to thyroid cell transformation has only been recently demonstrated. Thus Parma *et al.* (319a) found mutations in three of 11 autonomous (benign) thyroid nodules. One of these mutations was found at posi-

tion 623 (Ala → Ile) and the other two were found at position 619 (Asp → Gly). The mutations were associated with constitutively activated basal cyclase activity. Since screening has been linked to the third intracellular loop it is possible that more complete screening would locate mutations in other parts of the receptor (319b).

X. Tumor Suppressor Genes

The existence of tumor suppressor genes derived from three lines of evidence: 1) somatic cell hybridization of tumor with normal cells (320–323), 2) the realization of Knudson's 'two hit' postulate (324–326) in familial retinoblastoma (324), and 3) arising from the retinoblastoma paradigm (327–333), the correlation of loss of chromosomal segment heterozygosity with inactivation of putative tumor suppressor genes. The use of anonymous DNA markers has facilitated the search for the tumor loss of heterozygosity (LOH), since chromosomal loss extends to regions flanking suppressor genes (334, 335). Tumor suppressor genes may also be identified by difference cloning or differential display of mRNA (336, 337).

A. LOH in thyroid tumors

Studies have been undertaken to explore chromosomal structural and numerical abnormalities in thyroid tumors. The number of patients studied is, however, limited and has incriminated a variety of chromosomal regions (125; and Ref. 338 for review). Many may be secondary to thyroid cell transformation (338).

LOH involving chromosome 3p was implicated in and specific to follicular carcinoma (338). 3p LOH was described in small cell lung carcinoma (339), non-small cell lung carcinoma (340, 341), and breast and testicular tumors (342–344). Renal cell carcinoma harbors a more proximal deletion (3p14–21) (345) than small cell lung carcinoma (p21–23). It is intriguing that Matsuo *et al.* (346) reported LOH at 11q13 in approximately 14% of follicular adenomas. It is implied that some follicular adenomas will pursue a line of progression distinct from that in line to follicular carcinomas and thus subvert the assumption (3, 338) that follicular adenomas are intermediate between normal follicular cells and follicular carcinomas. Moreover, the chromosomal region implicated (346) harbors the multiple endocrine neoplasia I gene (347–349) as well as oncogenes and growth factor genes. The findings of LOH in thyroid tumors, therefore, bear confirmation in larger series of patients.

B. Retinoblastoma (Rb) gene

The Rb gene spans 180 Kb and 27 exons and maps to 13q14 (326, 350, 351). It encodes a 110-kDa nuclear phosphoprotein (350, 352). Rb binds the viral oncoproteins SV40 large T antigen, human adenovirus EA1, and papillavirus E7 (353–357). This binding inactivates Rb's growth suppressing function (358). All the viral oncoproteins bind to an underphosphorylated form of Rb (359).

The Rb protein switches between a hyperphosphorylated

and relatively underphosphorylated state in a cell cycle-specific manner (360–364). By binding them, pRb regulates the actions of the transcription factor E2F (365–370), and cyclins D₁ and D₃ (371, 372) which play critical roles in promoting cell cycle progression and parallel the already defined role of p107 (370–375). Seemingly, these factors bind in the same pRb pocket as do oncoproteins. Cyclin D₂, which accumulates in a cell cycle-dependent manner, is instrumental (in association with cd kinase 2) in the hyperphosphorylation of pRb and thus its release of cyclin D₁ and D₃ (372). This schema explains the negative regulation of *myc* expression by Rb/E2F complex and the observation that *myc* can overcome the ability of Rb but not p53 in affecting the G₁ phase arrest (376, 377). It also raises the possibility that mutations in or amplification of E2F or of specific cyclins may result in a phenotype similar to a loss of Rb (370, 371).


A subtle role of Rb in growth-inhibiting pathways is compatible with normal embryogenesis in familial retinoblastoma, where a defective gene is passed on in the germline and with the normal development of mice in which one allele has been inactivated (378–380). Mouse embryos in which both alleles have been disabled, however, fail to reach term and show abnormalities in neural and hematopoietic development (378–381).

Somatic Rb gene inactivation has now been implicated in a number of tumors (382), aside from retinoblastoma and sarcomas which are seen in relation to germline heterozygosity at the Rb locus (383, 384). The range of tumors involved is, however, likely to be more restricted than for mutations in p53. These include cervical carcinoma [whether or not they are positive for human papilloma virus sequences (385)], small cell and non-small cell lung carcinomas, bladder, and breast carcinomas (386–394). Some of these tumors also harbor p53 mutations (385).

The mechanisms that inactivate the Rb gene include deletions, insertions, or point mutations and have been uncovered in retinoblastomas, osteosarcoma, and soft tissue sarcoma which develop later in life in patients with retinoblastoma (350, 395, 396) as well as in a variety of tumors and tumor cell lines (382). Mutations that affect serine and threonine phosphorylation of Rb protein or its ability to bind oncoproteins have been mapped to two domains: codons 393–572 and codons 646–772 spread over exons 13–22 (397–399). The frequent deletions of exons 13–17 in retinoblastoma suggest that they may contain recombinational 'hot spots' (351). Exonic deletions of the Rb gene are usually accomplished by mutation at splice junction resulting in in-frame deletions (382, 385). Genomic DNA from such tumors may not exhibit obvious abnormalities (389, 400). These mutations result in lack of gene expression or a defective Rb protein (382, 401).

The development of thyroid tumors in transgenic mice in which SV40 T antigen was targeted to the thyroid gland under the influence of thyroglobulin promoter (402) led us to study Rb gene mutations in thyroid tumors. Mutant Rb alleles were found in 55% of thyroid carcinomas but in none of the benign tumors. The rates of Rb mutations were similar in anaplastic and differentiated thyroid carcinoma. The mutations were either attributable to abnormal Rb mRNA splic-

FIG. 6. Functional domains of the p53 protein. The activation domain of p53 has been mapped to the NH₂-end of the protein (residues 20–42). The heat shock protein (hsp) 70 binding domain is further upstream and overlaps the activation domain. The oligomerization domain (residues 344–393) is at the COOH end of p53, whereas the sequence necessary for nuclear localization is mapped to residues 316–325. Domains II–IV represent areas that are highly conserved among species. Some 98% of all p53 mutations described in tumors occur in these four domains (also see Fig. 7). *cdc2* Represents serine 315 phosphorylated by p34^{cdc2} kinase; CK2, serine 392 phosphorylated by casein kinase 2. [Adapted with permission from B. Vogelstein and K. W. Kinzler: *Cell* 70:523–526, 1992 (162). ©1992 by Cell Press.]



ing, C-terminal deletion, or point mutations (403). The most common mutations were deletions of exon 21, resulting in the fusion of exons 20–22. The deletion was, however, not present in genomic Rb gene sequences indicating that the deletion was the result of abnormal Rb mRNA splicing. One papillary carcinoma sample contained an in-frame deletion of most of exons 19 and 21 as well as the entire exon 20. The same sample also contained a deletion of exon 21. Twelve percent of malignant tumors harbored both Rb and p53 mutations. p53 And Rb double mutations appear to be more frequent in advanced disease stage. Thus Rb may be an important factor in thyroid cell malignant transformation, and possibly in tumor progression.

C. p53

The normal gene for p53 encodes a 393-residue long nuclear phosphoprotein mapping to chromosome 17p13 and spanning 11 exons. Exon 1 does not encode protein sequences (404). Cross-species comparison reveals five highly conserved domains, four of which fall within exons 5–8 (404): codons 117–142 (domain II); codons 171–181 (domain III); codons 234–258 (domain IV), and codons 270–286 (domain V). The C-terminal region of p53 was predicted to form an amphipathic helix-like structure (405, 406). p53 Oligomerizes through its C terminus to form tetramers (407). Its NH₂-terminus regulates the expression of downstream genes, which negatively control growth (408–412). The heat shock protein (hsp) 70 binding region (413) in the NH₂-terminus overlaps the activation domain. hsp 70 May act as a chaperone that facilitates oligomerization. The major nuclear localization signals are mapped to codons 316–325 (414) and include the target of serine phosphorylation by p34 *cd* kinase 2 (415). As most mutations impair the ability of p53 to bind DNA and/or result in dysfunction of the NH₂-terminus activation domains, they must affect the conformation of the entire protein (162, 416–418) (Fig. 6). Wild type p53, but none of the mutants tested, transactivates the putative promoter sequences (408). p53 Inhibits transcription from minimal promoters or it binds to TATA-binding protein, thus

inhibiting transcription indirectly (418–420).

p53 Appears to regulate growth-inhibiting signals at the G1-S transition of the cell cycle and monitor the genetic integrity at cell division in the same way as the *rad 9* gene in yeast (421). The effects of p53 are apparent only in stressed cells (162, 163, 422). Normal cells exposed to radiation (156), UV light, or UV-mimetic drugs (423) exhibit an increase in p53 expression and are arrested in G₁ until repair is effected. In contrast, cells harboring mutant p53 genes continue to divide and either die or accumulate genetic defects leading to tumorigenesis (157). In certain tumor systems, the prior activation of *myc* cooperates with *ras* in neoplastic transformation in the absence of p53 mutations, whereas in the absence of an activated gene there is a strong selection for p53 mutation by *myc* resulting in hyperplasia and not neoplasia (151, 424) and stresses the role of p53 in apoptosis (151).

p53 Was initially found through its association with SV40 large T antigen oncoprotein. Considered (425) an oncogene which cooperates, like *myc*, with *ras* in transforming cells (426, 427), it was later shown to be a tumor suppressor (147, 428). The p53 genes employed in those early experiments were inactivated by mutations (429). Like the Rb protein, p53 binds SV40 large T antigen, the E6 proteins of human papilloma virus, and EA1 of adenoviruses (425, 430–433). These oncogenic viruses sequester the p53 protein by binding it with p53 protein.

Germline p53 mutations are found in Li-Fraumeni syndrome (421), a familial cancer syndrome in which relatives develop diverse malignancies including breast carcinoma, sarcomas, and brain tumors (434, 435). Germline mutations have now been found in sporadic patients with cancers comprising Li-Fraumeni syndrome, specifically those with second tumors (436–439) and in patients with familial breast cancer (440, 441). p53 Mutations in Li-Fraumeni syndrome are widely distributed among the p53 genes between amino acid residues 72 and 325 (418, 421, 442, 443).

Somatic p53 mutations have been described in a wide variety of human cancers (428, 444–451). Some 98% of these mutations are found in that part of the molecule encoded by

FIG. 7. anapl Doma numb I, Ile; in a v tumo: 273 ir progr polyn

exon: ous t rare 454). 415), 175, gene: 135, comp or m (456' trans dons for n W ferer frequ trans cific hepa aflato pred in h muta dout (459) Lc is ch ever hum Inde




FIG. 7. Distribution of p53 mutations described in malignant thyroid tumors and in thyroid tumor lines. (▲, well differentiated tumors; ●, anaplastic tumors; ■, anaplastic cell lines; ○, well differentiated cell lines). II-IV represent domains highly conserved among species studied. Domain II includes residues 117-142; domain III, residues 171-181; domain IV, residues 234-258; and domain V, 270-286, the residues are numbered according to Soussi *et al.* (404). Single letter residue abbreviations are as follows: A, Ala; C, Cys; D, Asp; E, Glu; F, Phe; G, Gly; H, His; I, Ile; K, Lys; L, Leu; M, Met; N, Asn; P, Pro; Q, Glu; R, Arg; S, Ser; T, Thr; V, Val; W, Trp and Y, Try. Some 98% of all the mutations described in a variety of malignant tumors have been mapped to these four domains. By contrast, only 65% of the mutations described in malignant thyroid tumors or tumor lines map to these domains. Eleven of 13 mutations described in anaplastic tumors do map to domains III, IV, and V. Position 273 in homology domain V and 248 in domain IV appear to be mutation hot spots for anaplastic thyroid tumors and may be involved in tumor progression. It is highly unlikely that mutations outside the evolutionary conserved regions found in well differentiated tumors are rare polymorphisms.

exons 5-8 (428, 444). The missense mutations are so numerous that some are bound to represent previously undetected rare polymorphisms or may not affect p53 function (452-454). Mutated p53 proteins may bind DNA abnormally (408, 415), may exhibit NH₂-termini (e.g. mutants at codons 143, 175, 248, and 273) which are unable to activate reporter genes (408), may bind tightly to hsp 70 (mutants at codons 135, 175 but not 273) and to each other forming stable complexes and prolonging the mutant's half-life (429, 455), or may fail to bind to conformation-dependent antibodies (456) (mutants at codon 135, 175) or to suppress growth or transformation in transfection assays (162, 428). Three codons, 175, 248, and 273, appear to be particularly targeted for mutations (428).

When mutations are examined by tumor type, some differences emerge with respect to the position of the hot spots, frequency of mutation involving transitions compared to transversion (444). This exercise has yielded some site-specific information regarding the mutagens involved. Thus hepatocellular carcinoma in areas with high exposure to aflatoxin B1 ingestion and hepatitis B virus is associated predominantly with codon 249 mutation (457, 458). Likewise in human squamous cell carcinoma of the skin, p53 was mutated exclusively at pyrimidines, with frequent CC → TT double base changes, pathognomonic of UV light effect (459).

Loss of one allele with mutations in the remaining allele is characteristic of p53 inactivation (428). Unlike Rb, however, mutation at one p53 allele may also contribute to human carcinogenesis through negative dominant effects. Indeed, some mutant p53 proteins found in human tumors

can cooperate with the *ras* oncogene to transform primary cells, suggesting that they can inactivate endogenous wild-type p53 proteins (428). p53 Mutants differ in dominant negative effect, e.g. Mutants at codon 175 are severalfold more efficient than codon 273 mutants in cooperating with *ras* (428, 455). Wild-type p53 may also be bound and sequestered by E6 oncoprotein in human cervical carcinoma or the product of the MDM2 gene (162) found to be amplified in a high proportion of human sarcomas (459-461).

Mutations of the p53 gene in human neoplasia are generally regarded as late events (9, 462). There is, however, mounting evidence to show that it can also occur early in the neoplastic sequence in some malignancies (423, 446, 463, 464).

Mutations in the p53 gene appear to be relevant to the development and progression of malignant thyroid tumor. Initially, no mutations of p53 were detected in 129 thyroid tumor samples studied by immunohistochemistry or 20 tumor samples (465) by genomic DNA sequencing of amplified exons 5, 7, and 8. Both a cell line established from recurrent follicular carcinoma and the original tumor harbored a homozygous transitional (CGT → CAT) mutation at codon 273 (465). Concentration of mutations in exon 6 and/or gross contamination of tumor with nontumorous tissue has been invoked to account for the negative results in that study (416, 466).

Three other studies have documented p53 mutations in thyroid carcinoma (416, 466, 467). In two the mutations were predominantly or exclusively found in anaplastic tumors (466, 467), whereas in the third mutations were found in well differentiated malignant thyroid tumors (416). The geo-

TABLE 2. p53 Mutations in thyroid cancer

	Type	Stage	Age	Sex	Exon	Codon	Base change	Amino acid change
1.	AC	IV	46	M	5	159	GCC → CCG	Ala → Pro
						177	CCC → CC	1 bp deletion (frame shift)
2.	PC	IV	70	F	8	268	AAC → AGC	Asn → Ser
3.	FC	II	48	F	8	292	AAA → AGA	Lys → Arg
4.	PC	II	46	M	7	256	ACA → ACG	No
5.	PC	II	45	M	7	250	CCC → CTC	Pro → Leu
6.	PC	II	24	F	6	212	TTT → TTTT	1 bp addition (frame shift)
					8	282	CGG → TGG	Arg → Trp
7.	PC	I	23	F	8	283	CGC → CGT	No
8.	PC	II	25	F	7	231	ACC → GCC	Thr → Ala
9.	PC	I	32	F	6	222	GAG → GAA	No
10.	PC	I	36	F	8	268	AAC → CAC	Asn → His
11.	PC	I	26	F	7	237	ATG → ATA	Met → Ile
12.	PC	II	27	F	5	161	GCC → BCC	Ala → Thr

Review of histology of samples showed that samples 2, 7, and 8 had evidence of solid foci suggestive of dedifferentiation. AC, Anaplastic carcinoma; PC, papillary carcinoma; FC, follicular carcinoma.

[Reproduced with permission from Ref. 416. ©The Endocrine Society.]

graphic variation in the apparent role of p53 as a factor in tumor progression in the United States and Japan, and in tumor development in Saudi Arabia, is intriguing and may be related to genetic or environmental factors. Certainly, all three studies were reported from apparently iodide-sufficient areas. Scrutiny of the locations and nature of the mutations involved, most of which are transitional in nature, does not provide clues to potential environmental mutagens. In the studies by Ito *et al.* (467) and Zou *et al.* (416), the mutations appear to be relatively scattered in the p53 gene (Fig. 7). That all five p53 mutations detected among six anaplastic

carcinoma samples showed a CGT → CAT (Arg → His) transition at codon 273 in one study (466) is intriguing. Ninety percent of the point mutations in the anaplastic thyroid tumors and 50% in the well differentiated carcinoma (416) were G:C→A:T transitions. Positions 273 (466, 467) and 248 (467) (cpG dinucleotides) appear to be mutational hot spots in anaplastic thyroid carcinoma. Methylated cytosines occur exclusively at CpG dinucleotides and are a major site of spontaneous and probably enzyme-catalyzed deamination driven mutations (468, 469) of the p53 gene (468). The possibility that these mutational hot spots are preferen-




FIG. 8. The distribution of nm23 transcripts in thyroid tumors. There was no difference between benign nodular thyroid tissue samples (goiter), stages I through III of thyroid carcinoma. Samples from patients with anaplastic tumors and those at stage IV were significantly higher. These results suggest no correlation between the tumors' metastatic potential and nm23 transcript abundance. As close correspondence between transcript abundance and nm23 protein expression was described (507), the results cannot be explained away by variation in message stability. Thyroid tumors were staged on the basis of the TNM international classification (see Ref. 508). AC, Anaplastic carcinoma.

TABLE 3. Oncogene and tumor suppressor genes incriminated in thyroid tumors

Oncogene/antioncogene	Tumor type	Frequency (%)	Lesion	Environment
Ras	Adenomas	80	(H-ras 61) Point mutation	Iodide deficient
	Follicular carcinoma	50	(H-ras 61) Point mutation	Iodide deficient
	Adenomas	17	(H-ras 61) Point mutation	Iodide sufficient
	Follicular carcinoma	10	(H-ras 61) Point mutation	Iodide sufficient
	Anaplastic carcinoma	60	Point mutation	?
	Thyroid tumors ^a	60	(K-ras) Point mutation	Radiation
PTC/ret TPC	Papillary carcinoma	25	Rearrangement	?
Trk	Papillary carcinoma	10	Rearrangement	?
	Multinodular Goiters	5		
Met	Follicular carcinoma	22	Increased expression	?
	Papillary carcinoma	74		
	Poorly differentiated carcinoma	75		
c-myc	Malignant tumor ^b	57	Increased expression	?
	Papillary carcinoma	30		
	Adenoma	70		
c-fos	Malignant tumor ^b	60	Increased expression	
	Adenoma	90		
gsp	Differentiated carcinoma ^c	10	Point mutation	Iodide deficient
	Adenoma	3-25	Point mutation	
	Multinodular Goiter	5	Point mutation	
p53	Differentiated carcinoma ^c	25	Point mutation or Deletion or insertion	?
	Anaplastic carcinoma	86		
RB	Differentiated carcinoma ^c	54.5	Deletion or Mutation	?
	Anaplastic carcinoma	60		

^a Thyroid tumors: papillary, follicular, anaplastic, and adenoma.^b Malignant tumors: papillary, follicular, and anaplastic carcinoma.^c Differentiated carcinoma: papillary and follicular carcinoma.

tially used in different races or geographic locations requires further exploration (466, 467).

Eight of the 25 mutations including two in anaplastic thyroid carcinomas were outside the homology domains II-V (Fig. 7). As the mutations in well differentiated carcinomas do not coincide with any of the three polymorphisms described (453, 454, 470, 471) nor with nondeleterious mutation at codon 181 (452) and are restricted to malignant tumors (416), it is likely that they are causally related to malignant thyroid cell transformation. Their deleterious nature can, however, be definitively assessed by transfection assays.

Interestingly, all 20 mutations reported from tumor tissue involved only one p53 allele and cannot be attributed to contamination with normal thyroid tissues. By contrast, three of four thyroid carcinoma cell lines (465, 466) showed homozygous inactivation of p53, suggesting that this attribute may confer on thyroid epithelial cells the ability to grow in culture. Our data (416) suggest that the exon 5-8 domains of p53 are mutable in malignant thyroid tissue as well as the subject of frame shifts (Table 2). The reasons for these mutation pressures would be interesting to explore.

XI. Progression to Metastatic Potential

Two general and overlapping mechanisms may lead to the acquisition by cancer cells of metastatic potential: 1) malignant cells with a growth advantage in the primary lesions are more likely to metastasize (472); 2) cancer cells that metastasize acquire additional properties as a result of activation or inactivation of specific genes (473).

With stepwise accumulation of cellular events, malignant cells elaborate their own growth factors, become resistant to cytokines and growth factors that are normally inhibitory to normal cell growth, and may actually grow under the influence of these inhibitory factors. Primary tumors are thus clonally dominated by subpopulations of malignant cells with metastatic competence (474, 475). Since these cells frequently aberrantly manufacture their own growth factors and cytokines they develop a private autocrine loop that allows for multigrowth factor independence (476-478). Early in this process, the cells become growth factor competent by elaborating ectopically (479, 480) basic FGF (bFGF), NGF, PDGF, and IGF-1 (481, 482). Further malignant progression




FIG. 9. Current understanding of molecular events in thyroid carcinogenesis. The figure synthesizes proven and putative events as well as others of a speculative nature. The pathways leading to follicular or papillary carcinoma are divergent. The point(s) at which additional events select one of polyclonally proliferating cells for monoclonal growth and local dominance is unclear, and we do not know that hyperplasia is a prerequisite for microscopic papillary carcinoma foci. We are limited not only by lack of information at the precancerous stage but also by the small numbers of follicular carcinomas studied in most series. It is clear that *ras* oncogene activation is specific for the follicular carcinoma route and the rearrangement of *ret*-protooncogene and TRK1 protein tyrosine kinases for the papillary carcinoma fork. Seemingly, loss of heterozygosity (LOH) at chromosome 3p is specific for follicular carcinoma and implies the involvement of a tumor suppressor gene mapping to that chromosomal region. LOH at 11q13 may well be one of several mechanisms for clonal appearance of follicular adenomas. Because of the generally held role of tumor suppressor in tumor progression, we speculate that these adenomas have a high likelihood of becoming malignant. *p53* and *Rb* are assigned roles in malignant transformation of thyroid cells to papillary and probably follicular carcinoma and progression to anaplasia from follicular carcinoma. A role for *gsp* in follicular adenoma formation is secure. It is, however, possible that *gsp* mutations appear late in malignant thyroid tumor and perhaps determine tumor behavior.

in some epithelial cancers is associated with their transition from growth inhibition, e.g. by TGF β and IL-6 to growth stimulation by these factors (483–485). Sustainance of these highly malignant (and metastatic) cells by IL-6 has now been demonstrated in a variety of epithelial cancers (486–491). The relationship of the discrete multiple genetic events leading to malignancy to the increasing growth independence and subversion of inhibitory to stimulatory signals remains to be elucidated, as is the notion that resistance to one cytokine selects for resistance to other related or unrelated cytokines (472).

Given that beyond the size of 2–3 mm, solid tumors require their own blood supply, neurovascularization is a necessity for tumor success. These vessels also provide a means whereby cells with a high growth potential metastasize. Angiogenesis-promoting proteins synthesized by tumors include bFGF, EGF, vascular-endothelial growth factor, hepatocyte growth factor, pleiotrophin, and PDGF (285, 492, 493). The degree of angiogenesis determines tumor progress and its outcome (494, 495).

XII. nm23

A candidate for a specific gene product which influences metastatic potential of tumors is nm23 (473, 496). Nm23 was identified on the basis of its reduced steady state mRNA

levels in several tumor cell lines with high metastatic potential (497). The gene (NME1) (498) was later found to be highly homologous with *Drosophila* abnormal wing disc (awd) developmental gene (499) and with the nucleoside diphosphate (NDP) kinases (500–502). A second gene (nm23-H2) which is 88% identical to nm23-H1 has been described (503). The two isoforms of nm23, which map to 17q 21.3 (498, 504), correspond to A and B chains of NDP kinase (504, 505) and are assembled in variable combinations into hexamers (504).

The abundance of nm23 mRNA and protein showed a negative correlation with metastatic potential in experimental melanomas (497) and human breast cancer (506, 507). Furthermore, transfection of the nm23 gene into melanoma cells reduced their metastatic potential (509). Deletions or mutations of nm23 were found to correlate with metastatic behavior of tumors (510–512). The correlation of nm23 transcript number with metastases was, however, not noted in colonic cancer (513), neuroblastoma (514), and in some solid tumors (515).

We have studied the abundance of nm23 in 49 thyroid tumors with a wide range of histology. Our results indicate that nm23 transcript abundance did not negatively correlate with metastatic potential or with clinical staging of the disease in general. The highest level of nm23 transcripts was found in anaplastic carcinoma (Fig. 8). We have also used

single str
possibilit
and/or c
Lack of n
cDNA fr
and two
The nu
of two r
chains (c
different
and/or i
isotypes.
than the
preclude
participa
loss of c
It, of co
complex
propose
increase
NDP
tions in
that NE
ADP rit
clathrin
parently
change
anism w
behavior
of these
entiation
role for
recogni
tion fac

One
basis fo
reducin
crimina
sition u
is a mu.
may all
thyroid
other c
ences v

A nu
transfo
and/or
signals.
mediat
action
gates, i
lated a
recept
manne
growth
mation

single strand conformational polymorphism to explore the possibility that the nm23 gene may harbor somatic mutations and/or deletions in thyroid carcinoma. None were found. Lack of mutation was further confirmed by sequencing nm23 cDNA from four tumors, two with normal level expression and two with high expression (516).

The nm23 data should be interpreted in light of the identity of two nm23 gene products with the NDP kinase A and B chains (504, 505). The assembly of the two chains with different sets of intra- and intermolecular disulfide bonds and/or ratios of A and B chains could yield a number of isotypes. It appears that the A chain decreases much more than the B chain in metastatic tumors (507) and may thus preclude the formation of specific complexes in which its participation is critical. A more extreme example would be loss of copies of NME1 and NME2 by loss of heterozygosity. It, of course, remains to be seen as to which NDP kinase complex is actually involved in limiting metastases. This proposed mechanism would not, however, account for the increase in nm23 with dedifferentiation in some tumors.

NDP kinases may be responsible for high GTP concentrations in the proximity of G proteins (517). The suggestion that NDP kinase activates the small GTP binding protein, ADP ribosylation factor (518), which is necessary for non-clathrin coatamers, could not be supported (519, 520). Apparently Golgi membranes can specifically catalyze the exchange onto ADP ribosylation factor (519, 520). The mechanism whereby specific or all NDP kinases inhibit metastatic behavior is unclear, but is likely due to effects downstream of these gene products. That NDP kinase suppresses differentiation of leukemic cells in the mouse (521) suggested a role for nm23 in proliferation and differentiation. The recent recognition that nm23-H2 (but not nm23-H1) is a transcription factor for *myc* is intriguing (522).

XIII. Perspective

One aim of cancer research is to understand the etiological basis for carcinogenesis, with the long term objective of reducing cancer incidence. Epidemiological studies may incriminate environmental factors as well as genetic predisposition in the etiology of cancer. The recognition that cancer is a multistep process and the rapid elucidation of these steps may allow targeted prophylaxis and therapy. The picture in thyroid cancer is emerging at a modest pace compared to other cancers (Table 3). We have thus drawn from experiences with other cancers in this respect.

A number of cellular steps are necessary for thyroid cell transformation. These result in the delivery of constitutive and/or aberrant growth signals and loss of growth-inhibiting signals, thus disrupting the cell cycle. Growth signals can be mediated by oversecretion of growth factors, constitutive action of growth factor receptors, or their oncogene surrogates, aberrant expression of such receptors, or the unregulated action of transducer systems downstream from the receptors. Most of these cellular events act in a dominant manner. One or more genes involved in suppression of cell growth may have to be inactivated for malignant transformation or tumor progression to take place. The effects of

these genes are usually recessive but there are variations on this theme. Later, with tumor progression the transformed cells may subvert the action of growth-inhibiting cytokines and use them for growth. It is likely that early competence events lead to thyroid cell clonal growth. These cells then become the targets for the random accumulation of supplementary events, including inactivation of tumor suppressor genes which would lead to malignant transformation. It can be inferred that such malignant cells are programmed to alter expression of surface molecules including adhesion proteins as well as to induce proliferation of supporting connective tissue and capillaries which would dictate the biological behavior of these tumors. Figure 9 synthesizes both the defined and putative cellular genetic events involved in thyroid carcinogenesis.

*To see a world in a grain of sand
and a heaven in a wild flower
hold infinity in the palm of your hand
and eternity in an hour.*

William Blake, *Auguries of Innocence*

Acknowledgments

We wish to thank Ms. Colleen Bilkey and Ms. Ragenia Humphrey for painstaking preparation of the manuscript, Ms. Mabeth Martinez for help with the graphics, Mr. Phillip Ahring for preparing primers for many of the studies described, and Dr. Sultan Al-Sedairy for making available thyroid tumor tissue from the Tissue Bank for our studies.

We acknowledge the valuable contributions made by Dr. Valeria Stenzky and Ferenc Juhasz (Debrecen, Hungary), Desmond Robb, Allan Kwan, and Bodil Larsen (St John's, Newfoundland, Canada) to our earlier studies of thyroid cancer.

References

1. Scriver CR, Childs B 1989 Garrod's Inborn Factors in Disease. Oxford University Press, Oxford
2. Studer H, Peter HJ, Gerber H 1989 Natural heterogeneity of thyroid cells: the basis for understanding thyroid function and nodular goiter growth. *Endocr Rev* 10:125-135
3. Williams ED 1979 The aetiology of thyroid tumors. *Clin Endocrine Metab* 8:193-207
4. Foulds L 1958 The natural history of cancer. *J Chronic Dis* 8:2-37
5. Peto R 1977 Epidemiology multistage models, short term mutagenesis tests. In: *Origins of Human Cancer*. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, pp 1403-1428
6. Vogelstein B, Fearon ER, Kern EK, Hamilton SR, Preisinger AC, Nakamura Y, White R 1989 Allelotyping of colorectal carcinomas. *Science* 244:207-211
7. Kinzler KW, Nilbert MC, Su L-K, Vogelstein B, Bryan TM, Levy DB, Smith KJ, Preisinger AC, Hedge P, McKechnie D, Finnear R, Markham A, Groffen J, Boguski MS, Altschul SF, Horii A, Ando H, Miyoshi Y, Miki Y, Nishisho I, Nakamura Y 1991 Identification of FAP locus genes from chromosome 5q21. *Science* 253:661-665
8. Fearon ER, Cho KR, Nigro JM, Kern SE, Simons JW, Ruppert JM, Hamilton SR, Preisinger AC, Thomas G, Kinzler KW, Vogelstein B 1990 Identification of a chromosome 18q gene that is altered in colorectal cancers. *Science* 247:49-56
9. Fearon ER, Vogelstein B 1990 A genetic model for colorectal tumorigenesis. *Cell* 61:759-767
10. Kinzler KW, Nilbert MC, Su L-K, Vogelstein B, Bryan TM, Levy

- DB, Smith KJ, Preisinger AC, Hedge P, McKechnie D, Finnear R, Markham A, Groffen J, Boguski MS, Altschul SF, Horil A, Ando H, Miyoshi Y, Milki Y, Nishisho I, Nakamura Y 1991 Identification of a gene located at chromosome 5q21 that is mutated in colorectal cancers. *Science* 251:1366-1370
11. El-Deiry WS, Nelkin BD, Celano P, Yen RW, Falco JP, Hamilton SR, Baylin SB 1991 High expression of the DNA methyltransferase gene characterizes human neoplastic cells and progression stages of colon cancer. *Proc Natl Acad Sci USA* 88:3470-3474
 12. Powell SM, Zilz N, Beazer-Barclay Y, Bryan TM, Hamilton SR, Thibodeau SN, Vogelstein B, Kinzler KW 1992 APC mutations occur early during colorectal tumorigenesis. *Nature* 359:235-237
 13. Weinberg RA 1989 Oncogenes, antioncogenes and the molecular bases of multistep carcinogenesis. *Cancer Res* 49:3713-3721
 14. Zurzolo C, Gentile R, Mascia A, Garbi C, Polistina C, Aloj L, Avvedimento VE, Nitsch L 1991 The polarized epithelial phenotype is dominant in hybrids between polarized and unpolarized rat thyroid cell lines. *J Cell Sci* 98:65-73
 15. Munari-Silem Y, Mesnil M, Selmi S, Bernier-Valentin F, Rabilloud R, Rousset B 1990 Cell-cell interactions in the process of differentiation of thyroid epithelial cells into follicles: a study by micro injection and fluorescence microscopy on *in vitro* reconstituted thyroid follicles. *J Cell Physiol* 145:414-427
 16. Dunn JT 1991 Thyroglobulin: chemistry and biosynthesis. In: Braverman LE, Utiger RE (eds) *The Thyroid—Fundamental, Clinic Text*. JB Lippincott Co, Philadelphia, pp 98-110
 17. Dumont JE, Maenhaut C, Pirson I, Baptist M, Roger PP 1991 Growth factors controlling the thyroid gland. *Bailliere's Clin Endocrinol Metab* 5:727-754
 18. Coclet J, Foureau F, Ketelbant P, Galand P, Dumont JE 1989 All population kinetics in dog and human adult thyroid. *Clin Endocrinol (Oxf)* 31:655-665
 19. Conde E, Martin-Lacave I, Utrilla JC, Moreno A, Gonzales-Campora R, Galera-Davidson H 1992 Mitotic activity of the endocrine cells in rat thyroid glands during postnatal life. *Endocrinology* 131:436-440
 20. Dawson TP, Wyllie FS, Wynford-Thomas D 1991 *In vitro* responsiveness to serum growth factors is inversely related to *in vivo* malignancy in human thyroid epithelial cells. *Br J Cancer* 63:827-900
 21. Fayet G, Hovsepian S 1979 Demonstration of growth in porcine thyroid cell culture. *Biochimie* 61:923-930
 22. Gartner R, Greil W, Demharter P, Horn K 1985 Involvement of cyclic AMP, iodine and metabolites of arachidonic acid in the regulation of cell proliferation of isolated porcine thyroid follicles. *Mol Cell Endocrinol* 42:145-155
 23. Eggo MC, Bachrach LK, Fayet G, Errick J, Kudlow JE, Cohen MF, Burrow GN 1984 The effects of growth factors and serum on DNA synthesis and differentiation in thyroid cells in culture. *Mol Cell Endocrinol* 38:141-150
 24. Roger PP, Dumont JE 1984 Factors controlling proliferation and differentiation of canine thyroid cells cultured in reduced severe conditions: effects of thyrotropin, cyclic AMP and growth factors. *Mol Cell Endocrinol* 36:79-93
 25. Roger PP, Taton M, Van Sande J, Dumont JE 1988 Mitogenic effects of thyrotropin and cyclic AMP in differentiated human thyroid cells *in vitro*. *J Clin Endocrinol Metab* 66:1158-1165
 26. Williams DW, Williams ED, Wynford-Thomas D 1988 Loss of dependence on IGF-1 for proliferation of human thyroid adenoma cells. *Br J Cancer* 57:535-539
 27. Sun X, Rothenberg P, Kahn CR, Backer JM, Araki E, Wilden PA, Cahill DA, Goldstein BJ, Waite MF 1991 Structure of the insulin receptor substrate IRS-1 defines a unique signal transduction protein. *Nature* 352:73-77
 28. Condorelli G, Formisano P, Miele C, Beguinot F 1992 Thyrotropin regulates autophosphorylation and kinase activity in both the insulin and insulin-like growth factor-I receptors in FRTL5 cells. *Endocrinology* 130:1615-1621
 29. Roger PP, Dumont JE 1982 Epidermal growth factor controls the proliferation and the expression of differentiation in canine thyroid cells in primary culture. *FEBS Lett* 144:209-212
 30. Westermark K, Karlsson FA, Westermark B 1983 Epidermal growth factor modulates thyroid growth and function in culture. *Endocrinology* 112:1680-1686
 31. Lamy F, Taton M, Dumont JE, Roger PP 1990 Control of protein synthesis by thyrotropin and epidermal growth factor in human thyrocytes: role of morphological changes. *Mol Cell Endocrinol* 73:195-209
 32. Westermark K, Westermark B, Karlsson FA, Ericson LE 1986 Location of epidermal growth factor receptors on porcine-thyroid follicle cells and receptor regulation by thyrotropin. *Endocrinology* 118:1040-1046
 33. Atkinson S, Kendall-Taylor P 1987 Effect of thyrotropin on epidermal growth factor receptors in monolayer cultures of porcine thyroid cells. *J Endocrinol* 114:179-184
 34. Bachrach LK, Eggo MC, Mak WW, Burrow GN 1985 Phorbol esters stimulate growth and inhibit differentiation in cultured thyroid cells. *Endocrinology* 116:1603-1609
 35. Bachrach LK, Eggo MC, Hintz RL, Burrow GN 1988 Insulin-like growth factors in sheep thyroid cells: action, receptors and production. *Biochem Biophys Res Commun* 154:861-867
 36. Roger PP, Rouse S, Servais P, van Heuverswyn B, Dumont JE 1986 Stimulation of cell proliferation and inhibition of differentiation expression by tumor-promoting phorbol esters in dog thyroid cells in primary culture. *Cancer Res* 46:898-906
 37. Gerard CM, Roger PP, Dumont JE 1989 Thyroglobulin gene expression as a differentiation marker in primary cultures of calf thyroid cells. *Mol Cell Endocrinol* 61:23-35
 38. Minuto F, Barreca A, del Monte P, Carlola G, Torre GC, Giordano G 1989 Immunoreactive insulin-like growth factor I (IGF-I) and IGF-I-binding protein content in human thyroid tissue. *J Clin Endocrinol Metab* 68:621-626
 39. Maciel RMB, Moses AC, Villone G, Tramontano D, Ingbar SH 1988 Demonstration of the production and physiological role of insulin-like growth factor II in rat thyroid follicular cells in culture. *J Clin Invest* 82:1546-1553
 40. Greil W, Rafferzeder M, Brechtner G, Gartner R 1989 Release of an endothelial cell growth factor from cultured porcine thyroid follicles. *Mol Endocrinol* 3:858-865
 41. Rasmussen AK, Bech K, Feldt-Rasmussen U, Poulsen S, Siembaek-Nielsen K, Friis T, Bendtsen K 1987 The influence of interleukin-1 on the function of *in vitro* cultured human thyroid cells in monolayers. *Acta Endocrinol (Copenh)* 281:93-95
 42. Kawabe Y, Eguchi K, Shimomura Mine M, Otsubo T, Ueki Y, Tezuka H, Nakao H, Kawakami A, Migita K, Yamashita S, Matsunaga M, Ishikawa N, Ito K, Nagataki S 1989 Interleukin-1 production and action in thyroid tissue. *J Clin Endocrinol Metab* 68:1174-1183
 43. Weetman AP, Bennett GL, Wong WLT 1992 Thyroid follicular cells produce interleukin 8. *J Clin Endocrinol Metab* 75:328-330
 44. Pang XP, Park M, Hershman JM 1992 Transforming growth factor-beta blocks protein kinase-A-mediated iodide transport and protein kinase-C-mediated DNA synthesis in FRTL-5 rat thyroid cells. *Endocrinology* 131:45-50
 45. Grubeck-Loebenstein B, Buchan G, Sadeghi R, Kissonerghis M, Londei M, Turner M, Pirich K, Roka R, Nierderle B, Kassal H, Waldhausi W, Feldmann M 1989 Transforming growth factor beta regulates thyroid growth: role in the pathogenesis of non-toxic goiter. *J Clin Invest* 83:764-770
 46. Huber GK, Davies TF 1990 Human fetal thyroid growth *in vitro*: system characterization and cytokine inhibition. *Endocrinology* 126:869-875
 47. Yamashita S, Kimura H, Ashizawa K, Nagayama Y, Hirayu H, Izumi M, Nagataki S 1989 Interleukin-1 inhibits thyrotropin-induced human thyroglobulin gene expression. *J Endocrinol* 122:177-183
 48. Kimura H, Yamashita S, Namba H, Tominaga T, Tsuruta M, Yokoyama N, Izumi M, Nagataki S 1992 Interleukin 1 inhibits human carcinoma cell growth. *J Clin Endocrinol Metab* 75:596-602
 49. Goodman AL, Rone JD 1987 Thyroid angiogenesis: endothelial chemoattractant activity from rat thyroid cells in culture. *Endocrinology* 121:2131-2140
 50. Dumont JE, Jauniaux JC, Roger PP 1989 The cyclic AMP-me-

ture.
rotein
uman
rinol
1986
roid
ology
n on
roine
orbol
ured
-like
pro-
nt JE
ntia-
roid
gene
calf
Dior-
GF-I)
Clin
r SH
le of
ture.
lease
roid
iers-
e of
roid
di Y,
a S,
ikin-
etab
ular
30
with
and
roid
s M,
l H,
ctor
non-
itro:
logy
u H,
pin-
inol
i M,
ibits
196-
geli-
ure.
me-

- diated stimulation of cell proliferation. *Trends Biochem Sci* 14:67-71
51. Maenhaut C, Lefort A, Libert F, Parmentier M, Raspe E, Roger P, Corvillain B, Laurent E, Reuse S, Mockel J, Lamy F, Van Sande J, Dumont JE 1990 Function, proliferation and differentiation of the dog and human thyrocyte. *Horm Metab Res [Suppl]* 23:51-61
 52. Field JB, Ealey PA, Marshall NJ, Cockcroft S 1987 Thyroid-stimulating hormone stimulates increases in inositol phosphate as well as CAMP in the FRTL5 rat thyroid cell line. *Biochem J* 247:519-524
 53. Laurent E, Mockel J, Van Sande J, Graff I, Dumont JE 1987 Dual activation by thyrotropin of the phospholipase C and cyclic AMP cascades in human thyroid. *Mol Cell Endocrinol* 52:273-278
 54. Fujimoto J, Brenner-Gati L 1992 Protein kinase-C activation during thyrotropin stimulated proliferation of rat FRTL5 thyroid cells. *Endocrinology* 130:1587-1592
 55. Berridge MJ 1993 Inositol triphosphate and calcium signalling. *Nature* 361:315-325
 56. Raspe E, Reuse S, Roger PP, Dumont JE 1992 Lack of correlation between the activation of the Ca (2+) phosphatidylinositol cascade and the regulation of DNA synthesis in the dog thyrocyte. *Exp Cell Res* 198:17-26
 57. Van Sande J, Gremier G, Willems C, Dumont JE 1975 Inhibition by iodide of the thyroid cyclic 3', 5'-AMP system. *Endocrinology* 96:781-786
 58. Becks GP, Eggo MC, Burrow GN 1988 Organic iodide inhibits deoxyribo nucleic acid synthesis and growth in FRTL5 cells. *Endocrinology* 123:545-550
 59. Saji M, Kohn LD 1990 Effect of hydrocortisone on the ability of thyrotropin to increase deoxyribonucleic acid synthesis and iodide uptake in FRTL-5 rat thyroid cells: opposite regulation of adenosine 3',5' - monophosphate signal action. *Endocrinology* 127:1867-1876
 60. Dugrillon A, Bechtner G, Vedelhoven WM, Weber PC, Gartner R 1990 Evidence that iodolactone mediates the inhibitory effect of iodide on thyroid cell proliferation but not on adenosine-3',5' monophosphate. *Endocrinology* 127:337-343
 61. Pereira A, Braekman JC, Dumont JE, Boeynaems JM 1990 Identification of a major iodolipid from the horse thyroid gland as 2-iodohexadecanal. *J Biol Chem* 265:17018-17025
 62. Contor L, Lamy F, Lecocq R, Roger PP, Dumont JE 1988 Differential protein phosphorylation in induction of thyroid cell proliferation by thyrotropin epidermal growth factor or phorbol ester. *Mol Cell Biol* 8:2494-2503
 63. Lamy F, Roger PP, Lecocq R, Dumont JE 1989 Protein synthesis during induction of DNA replication in thyroid epithelial cells: evidence from late markers of distinct mitogenic pathways. *J Cell Physiol* 138:568-578
 64. Rossomando AJ, Payne DM, Weber MJ, Sturgill TW 1989 Evidence that pp42, a major tyrosine kinase target protein, is a mitogen activated serine/threonine protein kinase. *Proc Natl Acad Sci USA* 86:6940-6945
 65. Reuse S, Maenhaut C, Dumont JE 1990 Regulation of proto-oncogenes *c-fos* and *c-Myc* expression by protein tyrosine kinase, protein kinase c and cyclic AMP mitogenic pathways in dog primary thyrocytes: a positive and negative control by cyclic AMP on *c-Myc* expression. *Exp Cell Res* 189:33-40
 66. Heldin NE, Westermark B 1988 Epidermal growth factor, but not thyrotropin, stimulates the expression of *c-fos* and *c-myc* messenger ribonucleic acid in porcine thyroid follicle cells in primary culture. *Endocrinology* 122:1042-1046
 67. Heldin NE, Paulsson Y, Forsberg K, Heldin CH, Westermark B 1989 Induction of cyclic AMP synthesis by forskolin is followed by a reduction in the expression of *c-myc* messenger RNA and inhibition of ³H-thymidine incorporation in human fibroblasts. *J Cell Physiol* 138:17-23
 68. Office of Population Censuses and Surveys 1985. *Cancer Statistics Registrations for 1982, Series MBI: England and Wales*
 69. Biometry Branch, Division of Cancer Cause and Prevention 1975 Third national cancer survey incidence data. In: Cutler SJ, Young JL (eds) DHEW Publications No (NUT) 75-789, NCI Monograph No 41, Washington DC, p 25
 70. Silverberg E, Lubera TA 1989 *Cancer Statistics 1989*. CA 39:3-20
 71. Department of Health and Welfare 1959 The 2nd Fundamental Survey for Malignant Neoplasms. Tokyo, Japan
 72. Mubarak AL, Gjorgov AN, Ahmed J, Ismail AS 1986 Thyroid cancer in Kuwait: some epidemiological and clinical aspects. In: Khogali M, Omar YT, Gjorgov A, Ismail AS (eds) *Cancer Prevention in Developing Countries*. AS Pergamon Press, Oxford, UK, pp 2145-2150
 73. Henderson BE, Ross RK, Pike MC 1991 Toward the primary prevention of cancer. *Science* 254:1131-1138
 74. Ron E, Kleinerman RA, Boice Jr JD, Li Volsi VA, Flannery JT, Fraumeni JF 1987 A population-based case-control study of thyroid cancer. *J Natl Cancer Inst* 79:1-12
 75. American Cancer Society 1987 *Cancer Facts and Figures*. American Cancer Society Inc, New York
 76. Ries LAG, Hanakey BF, Miller BA, Hartman AM, Edwards BK 1991 *Cancer Statistics Review 1973-88*. National Cancer Institute, Bethesda, MD. NIH Publication No 91-2789
 77. Boring CC, Squires TS, Tong T 1993 *Cancer Statistics 1993*. CA: A Cancer Journal for Clinicians 43:7-26
 78. Hay ID 1990 Papillary thyroid carcinoma. *Clin Endocrinol Metab North Am* 19:545-576
 79. Moore Jr JH, Bacharach B, Choi HY 1985 Anaplastic transformation of metastatic follicular carcinoma of the thyroid. *J Surg Oncol* 29:216-221
 80. Venkatesh YS, Ordonez NG, Schultz PN, Hickey RC, Goepfert H, Samaan NA 1990 Anaplastic carcinoma of the thyroid: a clinicopathologic study of 121 cases. *Cancer* 66:321-330
 81. Heitz P, Moser H, Staub JJ 1976 Thyroid cancer. A study of 573 thyroid tumors and 161 autopsy cases observed over a thirty year period. *Cancer* 37:2329-2337
 82. Williams ED, Doniach I, Bjarnason O, Michie W 1977 Thyroid cancer in an iodide rich area: a histopathological study. *Cancer* 39:215-222
 83. Wahner HW, Cuello C, Correa P, Uribe LF, Gaitan E 1966 Thyroid carcinoma in an endemic goiter area, Cali, Colombia. *Am J Med* 40:58-66
 84. Franceschi S, Levi F, Negri E, Facina A, La Vecchio C 1991 Diet in thyroid cancer: a pooled analysis of four European case control studies. *Int J Cancer* 48:395-398
 - 84a. Belfiore A, La Rosa GL, La Porta GA, Giuffrida D, Milazzo G, Lupo L, Regalbuto C, Vigneri R 1992 Cancer risk in patients with cold thyroid nodules: relevance of iodine intake, sex, age and multinodularity. *Am J Med* 93:363-369
 85. Harach HR, Franssila KO, Wasenius VM 1985 Occult papillary carcinoma of the thyroid: a "normal" finding in Finland: a systematic autopsy study. *Cancer* 56:531-538
 86. Boice Jr JD, Engholm G, Kleinerman RA, Blettner M, Stovall M, Lisco H, Moloney WC, Austin DF, Bosch A, Cookfair DL, Krentz ET, Latourette HB, Merrill JA, Peters LJ, Schulz MD, Storm HH, Bjorkholm E, Pettersson F, Bell CM, Coleman MP, Fraser P, Neal FE, Prior P, Won Choi N, Hislop TG, Koch M, Kreiger N, Robb D, Robson D, Thomson DH, Lochmuller H, von Fournier D, Frischkorn R, Kjørstad KE, Rimpela A, Pejovic M-H, Kirm VP, Stankusova H, Berrino F, Sigurdsson K, Hutchison GB, MacMahon B 1988 Radiation dose and second cancer risk in patients treated for cancer of the cervix. *Radiat Res* 116:3-55
 87. Franceschi S, Fassina A, Talamini R, Mazzolini A, Vianello S, Bidoli E, Serraino D, La Vecchia C 1989 Risk factors for thyroid cancer in northern Italy. *Int J Epidemiol* 18:578-584
 88. Hamilton TE, van Belle G, LoGerfo JP 1987 Thyroid neoplasia in Marshall Islanders exposed to nuclear fallout. *JAMA* 258:629-636
 89. McTiernan AM, Weiss NS, Daling JR 1984 Incidence of thyroid cancer in women in relation to previous exposure to radiation therapy and history of thyroid disease. *J Natl Cancer Inst* 73:575-581
 90. Pottern LM, Kaplan MM, Larsen PR, Silva JE, Koenig RJ, Lubin JH, Stovall M, Boice Jr JD, 1990 Thyroid nodularity after child-

- hood irradiation for lymphoid hyperplasia: a comparison of questionnaire and clinical findings. *J Clin Epidemiol* 43:449-460
91. Prentice RL, Kato H, Yoshimoto K, Mason M 1982 Radiation exposure and thyroid cancer incidence among Hiroshima and Nagasaki residents. *Natl Cancer Inst Monogr* 62:207-212
 92. Preston-Martin S, Bernstein L, Pike MC, Maldonado AA, Henderson BE 1987 Thyroid cancer among young women related to prior thyroid disease and pregnancy history. *Br J Cancer* 55:191-195
 93. Rallison ML, Dobyns BM, Keating FR, Rall JE, Tyler FH 1975 Thyroid nodularity in children. *JAMA* 233:1069-1072
 94. Ron E, Modan B, Preston D, Alfandary E, Stovall M, Boice Jr JD 1989 Thyroid neoplasia following low-dose radiation in childhood. *Radiat Res* 120:516-531
 95. Schneider AB, Shore-Freedman E, Ryo UY, Bekerman C, Favus M, Pinsky S 1985 Radiation-induced tumors of the head and neck following childhood irradiation. *Medicine* 64:1-15
 96. Omran AR, Shore RE, Markoff RA, Friedhoff A, Albert RE, Barr H, Dahlstrom WG, Pasternack BS 1978 Follow-up study of patients treated by X-ray epilation for tinea capitis: psychiatric and psychometric evaluation. *Am J Pub Health* 68:561-567
 97. Shore RE, Woodard E, Hildreth N, Dvoretzky P, Hempelmann L, Pasternack B 1985 Thyroid tumors following thymus irradiation. *J Natl Cancer Inst* 74:1177-1184
 98. Wang JX, Boice Jr JD, Li BX, Zhang JY, Fraumeni JF 1988 Cancer among medical diagnostic x-ray workers in China. *J Natl Cancer Inst* 80:344-350
 99. Wang ZY, Boice Jr JD, Wei LX, Beebe GW, Zha YR, Kaplan MM, Tao ZF, Maxon HR III, Zhang SZ, Schneider AB, Tan B, Wesseler TA, Chen D, Ershow AG, Kleinerman RA, Littlefield LG, Preston D 1990 Thyroid nodularity and chromosome aberrations among women in areas of high background radiation in China. *J Natl Cancer Inst* 82:478-485
 100. Holm LE, Dahlqvist I, Israelsson A, Lundell G 1980 Malignant thyroid tumors after iodine-131 therapy. *N Engl J Med* 303:188-191
 101. Conard RA 1984 Late radiation effects in Marshall Islanders exposed to fallout 28 years ago. In: Boice Jr JD, Fraumeni JR (eds) *Radiation Carcinogenesis: Epidemiology and Biological Significance*. Raven Press, New York, pp 57-71
 102. Dobyns BM, Sheline GE, Workman JB, Tompkins EA, McConehney WM, Becker DV 1974 Malignant and benign neoplasms of the thyroid in patients treated for hyperthyroidism: a report of the cooperative thyrotoxicosis therapy follow-up study. *J Clin Endocrinol Metab* 38:976-978
 103. Wegelin C 1928 Malignant diseases of the thyroid gland and its relations to goiter in man and animals. *Cancer* 26:297-313
 104. Correa P, Llanos G 1966 Morbidity and mortality from cancer in Cali, Columbia. *J Natl Cancer Inst* 36:717-745
 105. Franssila K, Saxen E, Teppo L, Bjarnason O, Tulinius H, Normann T, Ringertz N 1981 Incidence of different morphological types of thyroid cancer in the Nordic countries. *Acta Pathol Microbiol Scand A* 89:49-55
 106. Pendergrast WJ, Milmore BK, Marcus SC 1961 Thyroid cancer and thyrotoxicosis in the United States: their relation to endemic goiter. *J Chronic Dis* 13:22-38
 107. Belfiore A, Garofalo MR, Giuffrida D, Runello F, Filetti S, Fiumara A, Ippolito O, Vigneri R 1990 Increased aggressiveness of thyroid cancer in patients with Graves' disease. *J Clin Endocrinol Metab* 70:830-835
 108. Behar R, Arganini M, Wu TC, McCormick M, Straus II FH, De Groot LJ, Kaplan EL 1986 Graves' disease and thyroid cancer. *Surgery* 100:1121-1127
 109. Farbota LM, Calandra DB, Lawrence AM, Paloyan E 1985 Thyroid carcinoma in Graves' disease. *Surgery* 98:1148-1153
 110. Hales IB, McElduff A, Crummer P, Clifton-Bligh P, Delbridge L, Hoschl R, Poole A, Reeve TS, Wilmshurst E, Wiseman J 1992 Does Graves' disease or thyrotoxicosis affect the prognosis of thyroid cancer. *J Clin Endocrinol Metab* 75:886-889
 111. Cady B 1991 Papillary carcinoma of the thyroid. *Semin Surg Oncol* 7:81-86
 112. Mazzaferrri EL 1991 Radioiodine and other treatments and outcomes. In: Braverman LE, Utiger RE (eds) *The Thyroid—A Fundamental Text*. JB Lippincott Co, Philadelphia, pp 1138-1165
 113. Tubiana M, Schlumberger M, Rougier P, Laplanche A, Benhamou E, Gardet P, Caillou B, Travagli JP, Parmentier C 1985 Long-term results and prognostic factors in patients with differentiated thyroid carcinoma. *Cancer* 55:794-804
 114. Samaan NA, Maheshwary YK, Naden S 1983 Impact of therapy for differentiated carcinoma of the thyroid: an analysis of 706 cases. *J Clin Endocrinol Metab* 56:1131-1137
 115. Plial RO, Bussey HJR, Thomson JPS 1987 Adenomatous polyposis: an association with carcinoma of the thyroid. *Br J Surg* 74:377-380
 116. Camiel MR, Mule JE, Alexander LI, Benninghoff DL 1968 Association of thyroid carcinoma with Gardner's syndrome in siblings. *N Engl J Med* 278:1056-1058
 117. Smith WG, Kern BB 1973 The nature of the mutation in familial multiple polyposis: papillary carcinoma of the thyroid brain tumors and familial multiple polyposis. *Dis Colon Rectum* 16:264-271
 118. Keshgegian AA, Enterline HT 1973 Gardner's syndrome duodenal adenomas, gastric adenomyoma and thyroid papillary follicular adenocarcinoma. *Dis Col Rectum* 21:255-260
 119. Hamilton SR, Bussey HJR, Mendelsohn G, Diamond MP, Pavlides G, Hutcheon D, Harbison M, Shermeta D, Morson BC, Yardley JH 1979 Ileal adenomas in nine patients with adenomatous polyposis coli/Gardner's syndrome. *Gastroenterology* 77:1252-1257
 120. Lee FI, MacKinnon ND 1981 Papillary thyroid carcinoma associated with polyposis coli. *Am J Gastroenterol* 76:138-140
 121. Thompson JS, Harned RK, Anderson JC, Hodgson PE 1983 Papillary carcinoma of the thyroid and familial polyposis coli. *Dis Colon Rectum* 26:583-585
 122. Sogol PB, Sugawara M, Gordon HE, Shellow WV, Hernandez F, Hershtman JW 1983 Cowden's disease: familial goiter and skin hamartomas. A report of three cases. *West J Med* 139:324-328
 123. Lote K, Andersen K, Nordal E, Brennhovd IO 1980 Familial occurrence of papillary thyroid carcinoma. *Cancer* 46:1291-1297
 124. Stoffer SS, Van Dyke DL, VadenBach J, Szpunar W, Weiss L 1986 Familial papillary carcinoma of the thyroid. *Am J Med Genet* 25:775-782
 125. Olah E, Balogh E, Bojan F, Juhasz F, Stenszky V, Farid NR 1990 Cytogenetic analysis of three papillary carcinomas and a follicular adenoma of the thyroid. *Cancer Genet Cytogenet* 44:119-129
 126. Hrafnkelisson J, Tulinius H, Jonasson JG, Oldsdottir G, Sigvaldason H 1989 Papillary thyroid carcinoma in Iceland: a study of the occurrence in families and the co-existence of other primary tumors. *Acta Oncol* 28:785-788
 127. Farid NR 1991 Genetic aspects of thyroid disease. In: Braverman LE, Utiger RE (eds) *The Thyroid—A Fundamental and Clinical Text*, ed 6. JB Lippincott Co, Philadelphia, pp 588-602
 128. Ozaki O, Ito K, Kobayashi K, Suzuki A, Manabe Y, Hosoda Y 1988 Familial differentiated, non-medullary thyroid carcinoma. *World J Surg* 112:565-571
 129. Cooper DS, Axelrod L, De Groot LJ, Vickery Jr AL, Maloof F 1981 Congenital goiter and the development of metastatic follicular carcinoma with evidence for a leak of non-hormonal iodide: clinical, pathological, kinetic and biochemical studies and a review of the literature. *J Clin Endocrinol Metab* 52:294-306
 130. McTiernan A, Weiss NS, Daling JR 1987 Incidence of thyroid cancer in women in relation to known or suspected risk factors for breast cancer. *Cancer Res* 1987 47:292-295
 131. Ron E, Curtis R, Hoffman DA, Flannery JT 1984 Multiple primary breast and thyroid cancer. *Br J Cancer* 49:87-92
 132. Tucker MA, Boice Jr JD, Hoffman DA 1985 Second cancer following cutaneous melanoma and cancers of the brain, thyroid, connective tissue, bone and eye in Connecticut 1935-1982. *Natl Cancer Inst Monogr* 68:161-89
 133. Schottenfeld D, Berg J 1971 Incidence of multiple primary cancers IV. Cancers of the female breast and genital organs. *J Natl Cancer Inst* 46:161-170
 134. Perkel VS, Gail MH, Lubin J, Pee DY, Weinstein R, Shore-Freedman E, Schneider AB 1988 Radiation-induced thyroid neo-

- plasmas: evidence for familial susceptibility factors. *J Clin Endocrinol Metab* 66:1316-1322
135. Larsen B, Thompson C, Kwan A, Farid NR 1986 Lack of association of HLA with thyroid cancer. *Tissue Antigens* 28:298-300
 136. Juhasz F, Stenszky V, Kozma L, Farid NR 1986 The relation of susceptibility to and biologic behavior of thyroid epithelial cell cancer to HLA-DR1. *Cancer* 58:52-54
 137. Juhasz F, Boros P, Szegedi G, Balazs G, Suranyi P, Kraszits E, Stenszky Farid NR 1989 Immunogenetic and immunologic studies of differentiated thyroid cancer. *Cancer* 63:1318-1326
 138. Sridama V, Hara Y, Fauchet R, De Groot LJ 1985 Association of differentiated thyroid carcinoma with HLA-DR7. *Cancer* 56:1086-1088
 139. Juhasz F, Balazs GY, Kozma L, Kraszits E, Stenszky V, Farid NR 1986 Interaction of IgG heavy-chain allotypes (Gm) and HLA in conferring susceptibility to thyroid carcinoma. *Clin Endocrinol (Oxf)* 25:17-21
 140. Melhem MF, Kunz HW, Gill III TJ 1993 A major histocompatibility complex-linked locus in rats critically influences resistance to diethylnitrosamine carcinogenesis. *Proc Natl Acad Sci USA* 90:1967-1971
 141. Dryja TP, Mukai S, Petersen R, Rapaport JM, Walton D, Yandell DW 1989 Parental origin of mutations of the retinoblastoma gene. *Nature* 339:556-558
 142. Toguchida J, Ishizaki K, Sasaki MS, Nakamura Y, Ikenaga M, Kato M, Sugimoto M, Kotoura Y, Yamamoto T 1989 Preferential mutation of paternally derived RB gene as the initial event in sporadic osteosarcoma. *Nature* 338:156-158
 143. Aaronson SA 1991 Growth factors and cancer. *Science* 254:1146-1153
 144. Solomon E, Borrow J, Goddard AD 1991 Chromosome aberrations and cancer. *Science* 254:1153-1160
 145. Matzuk MM, Finegold MJ, Su J-G, Hsueh AJW, Bradley A 1992 α -Inhibin is a tumor-suppressor gene with gonadal specificity in mice. *Nature* 360:313-319
 146. Clemens M 1992 Suppression with a difference. *Nature* 360:210-211
 147. Weinberg RA 1991 Tumor suppressor genes. *Science* 254:1138-1146
 148. Vaux DL, Cory S, Adams JM 1988 Bcl-2 gene promotes haemopoietic cell survival and cooperates with c-myc to immortalize pre-B cells. *Nature* 335:440-442
 149. Sentman CL, Shutter JR, Hockenbery D, Kanagawa O, Korsmeyer SJ 1991 bcl-2 inhibits multiple forms of apoptosis but not negative selection in thymocytes. *Cell* 67:879-888
 150. Garcia I, Martinov I, Tsujimoto Y, Martinov JC 1992 Prevention of programmed cell death of sympathetic neurons by the bcl-2 protooncogene. *Science* 258:302-304
 151. Marx J 1993 Cell death studies yield cancer clues. *Science* 259:760-761
 152. Bissonnette RP, Echeverri F, Mahdoubi A, Green DR 1992 Apoptotic cell death induced by c-Myc is inhibited by bcl-2. *Nature* 359:552-554
 153. Fanidi A, Harrington EA, Evan GI 1992 Cooperative interaction between c-Myc and bcl-2 proto-oncogene. *Nature* 359:554-556
 154. Kim SJ, Wagner S, Liu F, O'Reilly MA, Robbins PD, Green MR 1992. Retinoblastoma gene product activates expression of the human TGF-2 gene through transcription factor ATF-2. *Nature* 358:331-314
 155. Lin D, Shields MT, Ullrich SJ, Appella E, Mercer WE 1992 Growth arrest induced by wild-type p53 protein blocks cells prior to or near the restriction point in late G1 phase. *Proc Natl Acad Sci USA* 89:9210-9214
 156. Kastan MB, Zhan Q, el Deiry WS, Carrier F, Jacks T, Walsh WV, Plunkett BS, Vogelstein B, Fornace Jr AJ 1992 A mammalian cell cycle checkpoint pathway utilizing p53 and GADD45 is defective in ataxia-telangiectasia. *Cell* 71:587-597
 157. Hartwell L 1992 Defects in a cell cycle checkpoint may be responsible for the genomic instability of cancer cells. *Cell* 71:543-546
 158. Barrett JC, Tsutsumi T, Tlsty T, Oshimura M 1990 In: Harris CC, Liotta LA (eds) *Genetic Mechanisms in Carcinogenesis and Tumor Progression*. Wiley-Liss, Inc., New York, pp 97-114
 159. Hannan MA, Smith BP, Sigut D, Sackey K 1990 Radiosensitivity of fibroblasts obtained from a café-au-lait spot and normal appearing skin from a patient with neurofibromatosis (NF-6). *Cancer Genet Cytogenet* 47:191-196
 160. Hannan MA, Greer W, Smith BP, Sigut D, Ali MA, Amer MH 1991 Skin fibroblast cell lines derived from non-Hodgkin's-lymphoma (NHL) patients show increased sensitivity to chronic gamma irradiation. *Int J Cancer* 47:261-266
 161. Kuerbitz SV, Plunkett BS, Walsh WV, Kastan MB 1992 Wild type p53 is a cell cycle checkpoint determinant following irradiation. *Proc Natl Acad Sci USA* 89:7491-7495
 162. Vogelstein B, Kinzler KW 1992 p53 function and dysfunction. *Cell* 70:523-526
 163. Donehower LA, Harvey M, Slagle BL, McArthur MJ, Montgomery Jr CA, Butel JS, Bradley A 1992 Mice deficient for p53 are developmentally normal but susceptible to spontaneous tumors. *Nature* 356:215-221
 164. Goretzki PE, Lyons J, Stacy-Phipps S, Rosenau W, Demeure M, Clark OH, McCormick F, Rohrer HD, Bourne HR 1992 Mutational activation of RAS and GSP oncogenes in differentiated thyroid cancer and their biological implications. *World J Surg* 16:576-582
 165. Ishizaka Y, Kobayashi S, Ushijima T, Hirohashi S, Sugimura T, Nagao M 1991 Detection of ret TPC/PTC transcripts in thyroid adenomas and adenomatous goiter by an RT-PCR method. *Oncogene* 6:1667-1672
 166. Lyon MF 1988 X-chromosome inactivation and the location and expression of X-linked genes. *Am J Hum Genet* 42:8-16
 167. Vogelstein B, Fearon ER, Hamilton SR, Fairberg AP 1985 Use of restriction fragment length polymorphism to determine clonal origin of human tumors. *Science* 227:642-645
 168. Hicks DG, Li Volsi VA, Neidich JA, Puck JM, Kant JA 1990 Clonal analysis of solitary follicular nodules in the thyroid. *Am J Pathol* 137:553-562
 169. Thomas GA, Williams D, Williams ED 1989 The clonal origin of thyroid nodules and adenomas. *Am J Pathol* 134:141-147
 170. Namba H, Matsuo K, Fagin JA 1990 Clonal composition of benign and malignant human thyroid tumors. *J Clin Invest* 86:120-125
 171. Fey ME, Peter HJ, Hinds HL, Zimmermann A, Liechti-Gallati S, Gerber H, Studer H, Tobler A 1992 Clonal analysis of human tumors with M27 beta, a highly informative polymorphic X-chromosomal probe. *J Clin Invest* 89:1438-1444
 172. Namba H, Ross JL, Goodman D, Fagin JA 1992 Solitary polyclonal autonomous thyroid nodule: a rare cause of childhood hyperthyroidism. *J Clin Endocrinol Metab* 72:1108-1112
 173. Noguchi S, Motomura K, Inaji H, Imaoka S, Koyama H 1992 Clonal analysis of human breast cancer by means of the polymerase chain reaction. *Cancer Res* 52:6594-6597
 174. Gilliland DC, Blanchard K, Levy J, Perrin S, Bunn F 1991 Clonality in myeloproliferative disorder: analysis by means of the polymerase chain reaction. *Proc Natl Acad Sci USA* 88:6848-6852
 175. Frauman AG, Moses AC 1990 Oncogenes and growth factors in thyroid carcinogenesis. *Endocrinol Metab Clin North Am* 19:479-493
 176. Hashimoto T, Matsubara F, Mizukami Y, Miyazaki I, Michigishi T, Yanaihara N 1990 Tumor markers and oncogene expression in thyroid cancer using biochemical and immunohistochemical studies. *Endocrinol Jpn* 37:247-254
 177. Mizukami Y, Nonomura A, Hashimoto T, Michigishi T, Noguchi M, Matsubara F, Yanaihara N 1991 Immunohistochemical demonstration of epidermal growth factor and c-myc oncogene product in normal, benign and malignant thyroid tissues. *Histopathology* 18:11-18
 178. Di Carlo A, Mariano A, Pisano G, Parmeggiani V, Beguinot L, Macchia V 1990 Epidermal growth factor receptor and thyrotropin response in human tissue. *J Endocrinol Invest* 13:293-299
 179. Namba H, Gutman RA, Matsuo K, Alvarez A, Fagin JA 1990 H-ras proto-oncogene mutations in human thyroid neoplasms. *J Clin Endocrinol Metab* 71:223-229
 180. Aasland R, Lillehaug JR, Male R, Josendal O, Varhaug JE, Kleppe K 1988 Expression of oncogenes in thyroid tumours: coexpression of c-erbB2/neu and c-erbB. *Br J Cancer* 57:358-363

181. Lemoine NR, Wyllie FS, Lillehaug JR, Staddon SL, Hughes CM, Aasland R, Shaw G, Varhaug JE, Brown CL, Gullick WJ, Wynford-Thomas D 1990 Absence of abnormalities of the *c-erbB-1* and *c-erbB-2* proto-oncogenes in human thyroid neoplasia. *Eur J Cancer* 26:777-779
182. Sap J, Munoz A, Schmitt J, Stunnenberg H, Vennstrom B 1989 Repression of transcription mediated at a thyroid hormone response element by the *v-erbA* oncogene product. *Nature* 340:242-244
183. Damm K, Thompson CC, Evans RM 1989 Protein encoded by *v-erbB3A* functions as a thyroid-hormone receptor antagonist. *Nature* 339:593-597
184. Refetoff S, Weiss RE, Usala SJ 1993 The syndromes of resistance to thyroid hormone. *Endocr Rev* 14:348-399
185. Jasani B, Wyllie FS, Wright PA, Lemoine NR, Williams ED, Wynford-Thomas D 1990 Immunocytochemically detectable TGF-beta associated with malignancy in thyroid epithelial neoplasia. *Growth Factors* 2:149-155
186. Roberts AB, Sporn MB 1988 Transforming growth factor-B. *Adv Cancer Res* 51:107-145
187. Kroc M, Shandrasokar B, Yamanaki Y, Freiss H, Buchler M, Berger HG 1992 Overexpression of epidermal growth factor receptor in human pancreatic concern is associated with concomitant increases in the levels of epidermal growth factor and transforming growth factor alpha. *J Clin Invest* 90:1552-1560
188. Williams DW, Williams ED, Wynford-Thomas D 1989 Evidence for autocrine production of IGF-1 in human thyroid adenomas. *Mol Cell Endocrinol* 61:139-143
189. Heldin NE, Gustavsson B, Claesson-Welsh L, Hammacher A, Mark J, Heldin CH, Westermark B 1988 Aberrant expression of receptors for platelet-derived growth factor in an anaplastic thyroid carcinoma cell line. *Proc Natl Acad Sci USA* 85:9302-9306
190. Pawson T, Gish GD 1992 SH2 and SH3 domains from structure to function. *Cell* 71:359-362
191. Lavan BE, Kuhne MR, Garner CW, Anderson D, Reedijk M, Pawson T, Lienhard GE 1992 The association of insulin-elicited phosphotyrosine proteins with src homology 2 domains. *J Biol Chem* 267:11631-11636
192. Duchesne M, Schweighoffer F, Parker F, Clerc F, Frobert Y, Thang MN, Tocquet B 1993 Identification of SH3 domain of AP as an essential sequence for ras-GAP-mediated signalling. *Science* 259:525-528
193. Rozakis-Adcock M, McGlade J, Mbamalu G, Pelicci G, Daly R, Li W, Batzer A, Thomas S, Brugge J, Pelicci PG, Schlessinger J, Pawson T 1992 Association of the Shc and GRB2/Sem5 SH2-containing proteins is implicated in the activation of the Ras pathway by tyrosine kinases. *Nature* 360:689-692
194. McCormick F 1989 Ras GTPase activating protein: signal transmitter and signal terminator. *Cell* 56:5-8
195. McCormick F 1993 How receptors turn Ras on. *Nature* 363:15-16
196. Barbacid M 1987 ras genes. *Annu Rev Biochem* 56:779-827
197. Ballester R, Michaeli T, Ferguson K, Xu HP, McCormick F, Wigler M 1989 Genetic analysis of mammalian GAP expressed in yeast. *Cell* 59:681-686
198. Yatani A, Okabe K, Polakis P, Halenbeck R, McCormick F, Brown AM 1990 ras p21 And GAP inhibit coupling of muscarinic receptors to atrial K⁺ channels. *Cell* 61:769-776
199. Studer H, Gerber H, Zbaeren J, Peter H-J 1992 Histomorphological and immunohistochemical evidence that human nodular goiters grow by episodic replication of multiple clusters of thyroid follicular cells. *J Clin Endocrinol Metab* 75:1151-1158
200. Bos JL 1989 ras Oncogenes in human cancer: a review. *Cancer Res* 49:4682-4689
201. Almoguera C, Shibata D, Forrester K, Martin J, Arnheim N, Perucho M 1988 Most human carcinomas of the exocrine pancreas contain mutant c-K-ras gene. *Cell* 53:549-554
202. Rodenhuis S, Van de Wetering ML, Mooi WJ, Evers SG, van Zandwijk N, Bos JL 1987 Mutational activation of the K-ras oncogene, a possible pathogenetic factor in adenocarcinoma of the lung. *N Engl J Med* 317:929-935
203. Krengel U, Schlichting I, Scherer A, Schumann R, Frech M, John J, Kabsch W, Pai EF, Wittinghofer A 1990 Three dimensional structures of H-ras p21 mutants: molecular basis for their inability to function as signal switch molecules. *Cell* 62:539-548
204. Portella G, Ferulano G, Santoro M, Grieco M, Fusco A, Vecchio G 1989 The Kirsten murine sarcoma virus induces rat thyroid carcinomas *in vivo*. *Oncogene* 4:181-188
205. Lemoine NR, Staddon S, Bond J, Wyllie FS, Shaw JJ, Wynford-Thomas D 1990 Partial transformation of human thyroid epithelial cells by mutant Ha-ras oncogene. *Oncogene* 5:1833-1837
206. Shi YF, Zou MJ, Schmidt H, Juhasz F, Stenszky V, Robb D, Farid NR 1991 High rates of ras codon 61 mutation in thyroid tumors in an iodide-deficient area. *Cancer Res* 51:2690-2693
207. Bond J, Dawson T, Lemoine N, Wynford-Thomas D 1992 Effect of semi growth factors and phorbol ester on growth and survival of human thyroid epithelial cells expressing mutant ras. *Mol Carcinog* 5:129-135
208. Knyazev PG, Fedorov SN, Serova OM, Pluzhnikova GF, Novikov LB, Kalinovskiy VP, Seitz JF 1987 Molecular-genetic analysis of myc and c-Ha-ras proto-oncogene alterations in human carcinoma. *Hamatol Bluttransfus* 31:469-473
209. Knyazev PG, Nikiforova IF, Serova OM, Pluzhnikova GF 1990 Distribution and rearrangements of alleles of c-Ha-ras-1 protooncogene and their correlation with the development of lung, ovarian and thyroid cancers. *Neoplasma* 37:647-655
210. Papadimitriou K, Yiagnisis M, Tolis G, Spandidos DA 1988 Immunohistochemical analysis of the ras oncogene protein product in human thyroid neoplasms. *Anticancer Res* 8:1223-1227
211. Yoshida K, Hamatani K, Koide H, Ikeda H, Nakamura N, Akiyama H, Tsuchiyama H, Nakayama E, Shiku H 1988 Preparation of anti-ras M, 21,000 protein monoclonal antibodies and immunohistochemical analyses on expression of ras genes in human stomach and thyroid cancers. *Cancer Res* 48:5503-5509
212. Karga H, Lee JK, Vickery Jr AL, Thor A, Gaz RD, Jameson JL 1991 Ras oncogene mutations in benign and malignant thyroid neoplasms. *J Clin Endocrinol Metab* 73:832-836
213. Dockhorn-Dworniczak B, Caspari S, Schroder S, Bocker W, Dworniczak B 1990 [Demonstration of activated oncogenes of the ras family in human thyroid tumors using the polymerase chain reaction]. *Verh Dtsch Ges Pathol* 74:415-418
214. Namba H, Rubin SA, Fagin JA 1990 Point mutations of ras oncogenes are an early event in thyroid tumorigenesis. *Mol Endocrinol* 4:1474-1479
215. Schark C, Fulton N, Jacoby RF, Westbrook CA, Straus II FH, Kaplan EL 1990 N-ras 61 oncogene mutations in Hurthle cell tumors. *Surgery* 108:994-999
216. Suarez HG, du Villard JA, Severino M, Caillou B, Schlumberger M, Tubiana M, Parmentier C, Monier R 1990 Presence of mutations in all three ras genes in human thyroid tumors. *Oncogene* 5:565-570
217. Wright PA, Lemoine NR, Mayall ES, Wyllie FS, Hughes D, Williams ED, Wynford-Thomas D 1989 Papillary and follicular thyroid carcinomas show a different pattern of ras oncogene mutation. *Br J Cancer* 60:576-577
218. Lemoine NR, Mayall ES, Wyllie FS, Farr CJ, Hughes D, Padua RA, Thurston V, Williams ED, Wynford-Thomas D 1988 Activated ras oncogenes in human thyroid cancers. *Cancer Res* 48:4459-4463
219. Lemoine NR, Mayall ES, Williams ED, Thurston V, Wynford-Thomas D 1988 Agent-specific ras oncogene activation in rat thyroid tumors. *Oncogene* 3:541-544
220. Lemoine NR, Mayall ES, Wyllie FS, Williams ED, Goyns M, Stringer B, Wynford-Thomas D 1989 High frequency of ras oncogene activation in all stages of human thyroid tumorigenesis. *Oncogene* 4:159-164
221. Suarez HG, Du Villard JA, Caillou B, Schlumberger M, Tubiana M, Parmentier C, Monier R 1988 Detection of activated ras oncogenes in human thyroid carcinomas. *Oncogene* 2:403-406
222. Boultonwood J, Wyllie FS, Williams ED, Wynford-Thomas D 1988 N-myc expression in neoplasia of human thyroid C-cells. *Cancer Res* 48:4073-4077
223. Stringer BM, Rowson JM, Parkar MH, Seid JM, Hearn PR, Wynford-Thomas D, Ingemansson S, Woodhouse N, Goyns MH

- or their
9-548
Vecchio
thyroid
- ynford-
ithelial
- obb D,
thyroid
93
Effect
survival
of Car-
- Novi-
analysis
carci-
- F 1990
otoon-
varian
- A 1988
product
- Aki-
aration
muno-
stom-
- son JL
thyroid
- ker W,
of the
chain
- of ras
of En-
- II FH,
le cell
- berger
of mu-
ogene
- ies D,
licular
ogene
- Padua
Acti-
Res
- uford-
n rat
- is M,
f ras
nesis.
- iana
s on-
- 1988
ancer
- PR,
MH
- 1989 Detection of the H-RAS oncogene in human thyroid anaplastic carcinomas. *Experientia* 45:372-376
224. Wright PA, Williams ED, Lemoine NR, Wynford-Thomas D 1991 Radiation-associated and "spontaneous" human thyroid carcinomas show a different pattern of ras oncogene mutation. *Oncogene* 6:471-473
225. Roberts TM 1992 A signal chain of events. *Nature* 360:534-535
226. Kato GJ, Barrett J, Villa-Garcia M, Dang CV 1990 An amino-terminal c-myc domain required for neoplastic transformation activates transcription. *Mol Cell Biol* 10:5914-5920
227. Murre C, McCaw PS, Baltimore D 1989 A new DNA binding and dimerization motif in immunoglobulin enhancer binding, daughterless, MyoD, and myc proteins. *Cell* 56:777-783
228. Landschulz WH, Johnson PF, McKnight SL 1988 The leucine zipper: a hypothetical structure common to a new class of DNA binding proteins. *Science* 240:1759-1764
229. Amati B, Dalton S, Brooks MW, Littlewood TD, Evan GI, Land H 1992 Transcriptional activation by the human c-Myc oncoprotein in yeast requires interaction with Max. *Nature* 359:423-426
230. Kretzner L, Blackwood EM, Eisenman RN 1992 Myc and Max proteins possess distinct transcriptional activities. *Nature* 359:426-429
231. Amati B, Brooks MW, Levy N, Littlewood TD, Evan GI, Land H 1992 Oncogenic activity of the c-Myc protein requires dimerization with Max. *Cell* 72:233-245
232. Eilers M, Schirm S, Bishop JM 1991 The MYC protein activates transcription of the alpha-prothymosin gene. *EMBO J* 10:133-141
233. Evan GI, Wyllie AH, Gilbert CS, Littlewood TD, Land H, Brooks M, Waters CM, Penn LZ, Hancock DC 1992 Induction of apoptosis in fibroblasts by c-myc protein. *Cell* 69:119-128
234. Wyllie AH 1987 Apoptosis: cell death under homeostatic control. *Arch Toxicol [Suppl]* 11:3-10
235. Shi Y, Glynn JM, Guilbert LJ, Cotter TG, Bissonnette RP, Green DR 1990 Role for c-myc in activation-induced apoptotic cell death in T cell hybridomas. *Science* 257:212-218
236. Luscher B, Eisenman RN 1990 New light on Myc and Myb. I. *Myc. Genes Dev* 4:2025-2035
237. Kelly K, Siebenlist U 1986 The regulation and expression of c-Myc in normal and malignant cells. *Annu Rev Immunol* 4:317-318
238. del Senno L, Degli Uberti E, Rossi M, Buzzoni D, Barbieri R, Rossi P, Patracchini P, Bernardi F, Marchetti G, Conconi F, Gambiari R 1986 Identification of a c-myc oncogene lacking the exon 1 in the normal cells of a patient carrying a thyroid carcinoma. *FEBS Lett* 196:296-300
239. del Senno L, Gambiari R, degli Uberti E, Barbieri R, Bernardi F, Buzzoni D, Marchetti G, Pansini G, Perrotta C, Conconi F 1987 c-myc oncogene alterations in human thyroid carcinomas. *Cancer Detect Prev* 10:159-166
240. Terrier P, Sheng ZM, Schlumberger M, Tubiana M, Caillou B, Travagli JP, Fragu P, Parmentier C, Riou G 1988 Structure and expression of c-myc and c-fos proto-oncogenes in thyroid carcinomas. *Br J Cancer* 57:43-47
241. Burman KD, Djuh YY, La Rocca RV, Nunes ME, D'Avis JC, Nicholson DE, Wartofsky L 1987 c-myc expression in the thyroid. I. Normal, adenomatous and cancerous thyroid tissue. *Horm Metab Res [Suppl]* 17:63-65
242. Wyllie FS, Lemoine NR, Williams ED, Wynford-Thomas D 1989 Structure and expression of nuclear oncogenes in multi-stage thyroid tumorigenesis. *Br J Cancer* 60:561-565
243. Yamashita S, Ong J, Fagin JA, Melmed S 1986 Expression of the myc cellular proto-oncogene in human thyroid tissue. *J Clin Endocrinol Metab* 63:1170-1173
244. Shi Y, Zou M, Farid NR 1993 Expression of thyrotropin receptor gene in thyroid carcinoma is associated with good prognosis. *Clin Endocrinol (Oxf)* 39:269-274
245. Brabant G, Maenhaut C, Kohrle J, Scheumann G, Dralle H, Hoang-Vu C, Hesch RD, von zur Muhlen A, Vassart G, Dumont JE 1991 Human thyrotropin receptor gene: expression in thyroid tumors and correlation to markers of thyroid differentiation and dedifferentiation. *Mol Cell Endocrinol* 82:R7-12
246. Curran T 1988 The fos oncogene. In: Reddy EP, Skalka AM, Curran T (eds) *The Oncogene Handbook*. Elsevier, New York, pp 307-325
247. Lamph WW, Wamsley P, Sassone-Corsi P, Verma IM 1988 Induction of proto-oncogene *jun/AP-1* by serum and TPA. *Nature* 334:629-631
248. Nakamura T, Datta R, Sherman ML, Kufe D 1990 Regulation of c-jun gene expression by cAMP in HL-60 myeloid leukemia cells. *J Biol Chem* 265:22011-22015
249. Sassone-Corsi P, Visvader J, Ferland L, Mellon PL, Verma IM 1988 Induction of proto-oncogene fos transcription through the adenylate cyclase pathway: characterization of a cAMP-responsive element. *Gene Dev* 2:2529-2538
250. Auwerx J, Staels B, Sassone-Corsi P 1990 Coupled and uncoupled induction of fos and jun transcription by different second messengers in cells of hematopoietic origin. *Nucleic Acids Res* 18:221-228
251. Sassone-Corsi P, Ransone LJ, Lamph WW, Verma IM 1988 Direct interaction between fos and jun nuclear oncoproteins: role of the "leucine zipper" domain. *Nature* 336:692-695
252. Grieco M, Santoro M, Berlingieri MT, Melillo RM, Donghi R, Bongarzoni I, Pierotti MA, Della Porta G, Fusco A, Vecchio GP 1990 PTC is a novel rearranged form of the ret proto-oncogene and is frequently detected *in vivo* in human thyroid papillary carcinomas. *Cell* 60:557-563
253. Grieco M, Santoro M, Berlingieri MT, Donghi R, Pierotti MA, Della Porta G, Fusco A, Vecchio G 1988 Molecular cloning of PTC, a new oncogene found activated in human thyroid papillary carcinomas and their lymph node metastases. *Ann NY Acad Sci* 551:380-381
254. Ishizaka Y, Tahira T, Ochiai M, Ikeda I, Sugimura T, Nagao M 1988 Molecular cloning and characterization of human ret-II oncogene. *Oncogene Res* 3:193-197
255. Ishizaka Y, Ushijima T, Sugimura T, Nagao M 1990 cDNA cloning and characterization of ret activated in a human papillary thyroid carcinoma cell line. *Biochem Biophys Res Commun* 168:402-408
256. Takahashi M, Ritz J, Cooper GM 1985 Activation of a novel human transforming gene, ret, by DNA re-arrangement. *Cell* 42:581-588
257. Ishizaka Y, Itoh T, Tahira I, Ikeda T, Sugimura J, Tucker A, Fertitta A, Carrano AV, Nagao M 1989 Human ret proto-oncogene is mapped to chromosome 10q11.2. *Oncogene* 4:1519-1521
258. Donghi R, Sozzi G, Pierotti MA, Biunno I, Miozzo M, Fusco A, Grieco M, Santoro M, Vecchio G, Spurr NK, Della Porta G 1989 The oncogene associated with human papillary thyroid carcinoma (PTC) is assigned to chromosome 10 q11-q12 in the same region as multiple endocrine neoplasia type 2A (MEN2A). *Oncogene* 4:521-523
259. Sozzi G, Pierotti MA, Miozzo M, Donghi R, Radice P, De Benedetti V, Grieco M, Santoro M, Fusco A, Vecchio G, Mathew CGP, Ponder BAJ, Spurr NK, Della Porta G 1991 Refined localization to contiguous regions on chromosome 10q of the two genes (H4 and RET) that form the oncogenic sequence PTC. *Oncogene* 6:339-342
260. Fusco A, Grieco M, Santoro M, Berlingieri MT, Pilotti S, Pierotti MA, Della Porta G, Vecchio G 1987 A new oncogene in human thyroid papillary carcinomas and their lymph-nodal metastases. *Nature* 328:170-172
261. Jhiang SM, Caruso DR, Gilmore E, Ishizaka Y, Tahira T, Nagao M, Chiu I-M, Mazzaferri EL 1992 Detection of the PTC/ret^{PTC} oncogene in human thyroid cancer. *Oncogene* 7:1331-1337
262. Pierotti MA, Santoro M, Jenkins RB, Sozzi G, Bongarzoni I, Grieco M, Monzini N, Miozzo M, Herrmann MA, Fusco A, Hay ID, Della Porta G, Vecchio G 1992 Characterization of an inversion on the long arm of chromosome 10 juxtaposing D10S170 and RET and creating the oncogenic sequence RET/PTC. *Proc Natl Acad Sci USA* 89:1616-1620
263. Tahira T, Ishizaka Y, Itoh F, Nakayasu M, Sugimura T, Nagao M 1991 Expression of the ret proto-oncogene in human neuroblastoma cell lines and its increase during neuronal differentiation induced by retinoic acid. *Oncogene* 6:2333-2338
264. Santoro M, Rosati R, Grieco M, Berlingieri MT, D'Amato GL,

- de Franciscis V, Fusco A 1990 The ret proto-oncogene is consistently expressed in human pheochromocytomas and thyroid medullary carcinomas. *Oncogene* 5:1595-1598
265. Ikeda I, Ishizaka Y, Tahira T, Suzuki T, Onda M, Sugimura T, Nagao M 1990 Specific expression of the ret proto-oncogene in human neuroblastoma cell lines. *Oncogene* 5:1291-1296
266. Lanzi C, Borrello MG, Bongarzone I, Migliazza A, Fusco A, Grieco M, Santoro M, Gambeta RA, Zumima F, Della Porta G, Pierotti MA 1992 Identification of the product of two oncogenic re-arranged forms of the RET proto-oncogene in papillary thyroid carcinomas. *Oncogene* 7:1189-1194
267. Ishizaka Y, Ochiai M, Tahira T, Sugimura T, Nagao M 1989 Activation of ret II oncogene without a sequence encoding a transmembrane domain and transforming activity of two ret-II oncogene products differing in carboxy-termini due to alternative splicing. *Oncogene* 4:789-794
268. Takahashi M, Buma Y, Iwamoto T, Inaguma Y, Ikeda M, Hiai H 1989 Cloning and expression of ret proto-oncogene encoding a tyrosine kinase with two potential transmembrane domains. *Oncogene* 3:571-578
269. Takahashi M, Buma Y, Taniguchi M 1991 Identification of the ret proto-oncogene products in neuroblastoma and leukemia cells. *Oncogene* 6:297-301
270. Takahashi M, Buma Y, Hiai Y 1989 Isolation of ret proto-oncogene cDNA with amino-terminal signal sequence. *Oncogene* 4:805-806
271. Santoro M, Carlomagno F, Hay ID, Herrmann MA, Grieco M, Melillo R, Pierotti MA, Bongarzone I, Della Porta G, Berger N, Peix JL, Paulin C, Fabien N, Vecchio G, Jenkins RB, Fusco A 1992 Ret oncogene activation in human thyroid neoplasms is restricted to the papillary cancer subtype. *J Clin Invest* 89:1517-1522
272. Zou MJ, Shi YF, Farid NR 1994 Low rate of ret proto-oncogene activation (PTC/ret^{PTC}) in papillary thyroid carcinomas from Saudi Arabia. *Cancer* 73:176-180
273. Mulligan LM, Kwok JB, Healey CS, Elsdon MJ, Eng C, Gardner E, Love DR, Mole SE, Moore JJK, Pap L, Ponder MA, Telenius H, Tunnacliffe A, Ponder BAJ 1993 Germ-line mutations of the RET proto-oncogene in multiple endocrine neoplasia type 2A. *Nature* 363:458-460
274. Jhiang SM, Caruso DR, Mazzaferrri GL 1991 Detection of the PTC oncogene in human thyroid cancers. *Clin Res* 39:376A
275. Fabien N, Paulin C, Santoro M, Berger N, Grieco M, Galvain D, Barbier Y, Dubois PM, Fusco A 1992 Detection of RET oncogene activation in human papillary carcinoma by *in situ* hybridization. *Br J Cancer* 66:1094-1098
276. Santoro M, Melillo KM, Grieco M, Berlingieri MT, Vecchio G, Fusco A 1993 The TRK and tyrosine kinase oncogenes cooperate with ras in the neoplastic transformation of a rat ligand epithelial line. *Cell Growth Differentiation* 4:77-84
277. Bongarzone I, Pierotti MA, Monzini N, Mondellini P, Manenti G, Donghi R, Pilotti S, Grieco M, Santoro M, Fusco A, Vecchio G, Della Porta G 1989 High frequency of activation of tyrosine kinase oncogenes in human papillary thyroid carcinoma. *Oncogene* 4:1457-1462
278. Morris CM, Hao QL, Heisterkamp N, Fitzgerald PH, Groffen J 1991 Localization of the TRK proto-oncogene to human chromosome bands 1q23-1q24. *Oncogene* 6:1093-1095
279. Hempstead BL, Martin-Zanca D, Kaplan DR, Parada LF, Chao MV 1991 High-affinity NGF binding requires coexpression of the trk proto-oncogene and the low-affinity NGF receptor. *Nature* 350:678-683
280. Greco A, Pierotti MA, Bongarzone I, Pagliardini S, Lanzi C, Della Porta G 1992 TRK-T1 is a novel oncogene formed by the fusion of TPR and TRK genes in human papillary thyroid carcinomas. *Oncogene* 7:237-242
281. Comoglio PM 1993 Structure, biosynthesis and biochemical properties of the HGF receptor in normal and malignant cells. *Exper Suppl (Basel)* 65:131-165
282. Naldini L, Tamagone L, Vigna E, Sachs M, Hartmann G, Birchmeier W, Daikuhara Y, Tsubouchi H, Blasi F, Comoglio PM 1992 Extracellular proteolytic cleavage by urokinase is required for activation of hepatocyte growth factor/scatter factor. *EMBO J* 11:4825-4833
283. Rodrigues GA, Park M 1993 Isoforms of the met receptor tyrosine kinase. *Exper Suppl (Basel)* 65:167-179
284. Weidner KM, Hartmann G, Naldini L, Comoglio PM, Sachs M, Fonatsch C, Rieder H, Birchmeier W 1993 Molecular characteristics of HGF-SF and its role in cell motility and invasion. *Exper Suppl (Basel)* 65:322-328
285. Tajima M, Matsumoto K, Nakamura T 1992 Regulation of cell growth and motility by hepatocyte growth factor and receptor expression in various cell species. *Exp Cell Res* 202:423-431
286. Matsumoto K, Nakamura J 1992 Hepatocyte growth factor: molecular structure, roles in liver regeneration and other biological functions. *Crit Rev Oncog* 3:27-54
287. Di Renzo MF, Narsimhan RP, Olivero M, Bretti S, Giordano S, Medico E, Gaglia P, Zara P, Comoglio PM 1991 Expression of the Met/HGF receptor in normal and neoplastic human tissues. *Oncogene* 6:1977-2003
288. Di Renzo MF, Olivero M, Ferro S, Prat M, Bongarzone I, Pilotti S, Belfiore A, Costantino A, Vigneri R, Pierotti MA, Comoglio PM 1992 Overexpression of the c-MET/HGF receptor gene in human thyroid carcinomas. *Oncogene* 7:2549-2553
289. Zarnegar R, Muga S, Rahija R, Michalopoulos G 1990 Tissue distribution of hepatopoietin-A: a heparin-binding polypeptide growth factor for hepatocytes. *Proc Natl Acad Sci USA* 87:1252-1256
290. Bourne HR, Sanders DA, McCormick F 1991 The GTPase superfamily: conserved structure and molecular mechanisms. *Nature* 349:117-127
291. Kaziro Y, Itoh H, Kozasa T, Nakafuku M, Satoh T 1991 Structure and function of signal-transducing GTP binding proteins. *Annu Rev Biochem* 60:349-400
292. Spiegel AM, Shenker A, Weinstein LS 1992 Receptor-effector coupling by G proteins: implications of normal and abnormal signal transduction. *Endocr Rev* 13:536-565
293. Birnbaumer L 1992 Receptor to effector signalling through G proteins: roles for $\beta\gamma$ dimers as well as α subunits. *Cell* 71:1069-1072
294. Weinstein LS, Shenker A, Gejman PV, Merino MJ, Friedman E, Spiegel AM 1991 Activating mutations of the stimulatory G protein in the McCune-Albright syndrome. *N Engl J Med* 325:1688-1695
295. Schwindinger WF, Francomano CA, Levine MA 1992 Identification of a mutation in the gene encoding the α subunit of the stimulatory G protein of adenylyl cyclase in McCune-Albright syndrome. *Proc Natl Acad Sci USA* 89:5152-5156
296. Weinstein LS, Gejman PV, Friedman E, Kadowaki T, Collins RM, Gershon ES, Spiegel AM 1990 Mutations of the Gs α -subunit gene in Albright hereditary osteodystrophy detected by denaturing gradient gel electrophoresis. *Proc Natl Acad Sci USA* 87:8287-8290
297. Patten JL, Johns DR, Valle D, Eil C, Gruppuso PA, Steele G, Smallwood PM, Levine MA 1990 Mutation in the gene encoding the stimulatory G protein of adenylyl cyclase in Albright's hereditary osteodystrophy. *N Engl J Med* 322:1412-1419
298. Landis CA, Masters SB, Spada A, Pace AM, Bourne HR, Vallar L 1989 GTPase inhibiting mutations activate the α chain of Gs and stimulate adenylyl cyclase in human pituitary tumors. *Nature* 340:692-696
299. Landis CA, Harsh G, Lyons J, Davis RL, McCormick F, Bourne HR 1990 Clinical characteristics of acromegalic patients whose pituitary tumors contain mutant Gs protein. *J Clin Endocrinol Metab* 71:1416-1420
300. Lyons J, Landis CA, Harsh G, Vallar L, Grunewald K, Feichtinger H, Duh Q-Y, Clark OH, Kawasaki E, Bourne HR, McCormick F 1990 Two G protein oncogenes in human endocrine tumors. *Science* 249:655-659
301. Clementi E, Margaretti N, Meldolesi J, Taramelli R 1990 A new constitutively activating mutation of the Gs protein α subunit-gp oncogene is found in human pituitary tumors. *Oncogene* 5:1059-1061
302. Suarez HG, du Villard JA, Caillou B, Schlumberger M, Farmer

- BO J tier C, Monier R 1991 GSP mutations in human thyroid tumours. *Oncogene* 6:677-679
- osine 303. O'Sullivan C, Barton CM, Staddon SL, Brown CL, Lemoine NR 1991 Activating point mutations of the gsp oncogene in human thyroid adenomas. *Mol Carcinogen* 4:345-349
- as M, 304. Matuso K, Freidman E, Gejman P, Fagin JA 1993 The thyrotropin receptor (TSH-R) is not an oncogene for thyroid tumors and the α subunit of Gs in human thyroid neoplasms. *J Clin Endocrinol Metab* 76:1446-1451
- acter- 305. Osawa S, Dhanasekaran N, Woon CW, Johnson GL 1990 $G_{\alpha s}$ chimeras define the function of α chain domains in control of G protein activation and $\beta\gamma$ subunit complex interactions. *Cell* 63:697-706
- Exper 306. Maenhaut C, Van Sande J, Libert F, Abramowicz M, Parmentier M, Venderhaegen JJ, Dumont JE, Vassart G, Schiffman S 1990 RDC8 codes for an adenosine A_2 receptor with physiological constitutive activity. *Biochem Biophys Res Commun* 173:1169-1178
- of cell 307. Ledent C, Dumont FE, Vassart G, Parmentier M 1992 Thyroid expression of an A_2 adenosine receptor transgene induces thyroid hyperplasia and hyperthyroidism. *EMBO J* 11:537-542
- eptor 308. Selzer E, Wilfring A, Schiferer A, Herman M, Grubeck-Loebenstein B, Freissmuth M 1993 Stimulation of human thyroid growth via inhibitory guanine nucleotide binding (G) protein G_i : constitutive expression of the G-protein α subunit $G_{i\alpha-1}$ in autonomous adenoma. *Proc Natl Acad Sci USA* 90:1609-1613
- mo- 309. Vassart G, Dumont JE 1992 The thyrotropin receptor and the regulation of thyrocyte function and growth. *Endocr Rev* 13:596-611
- ogical 310. Chazenbalk GD, Nagayama Y, Russo D, Wadsworth HL, Rapoport B 1990 Functional analysis of the cytoplasmic domains of the human thyrotropin receptor by site-directed mutagenesis. *J Biol Chem* 265:20970-20975
- no S, 311. Nagayama Y, Wadsworth HL, Chazenbalk GD, Russo D, Seto P, Rapoport B 1991 Thyrotropin-luteinizing hormone/chorionic gonadotropin receptor extracellular domain chimeras as probes for TSH receptor function. *Proc Natl Acad Sci USA* 88:902-905
- on of 312. Mishari M, Loosfelt H, Atger H, Sar S, Guiochon-Mantel A, Milgrom E 1990 Cloning sequencing and expression of human TSH receptor. *Biochem Biophys Res Commun* 166:394-403
- issues. 313. Graves PN, Tomer Y, Davies TF 1992 Cloning and sequencing of a 1.3 Kb variant of the human thyrotropin receptor mRNA lacking the transmembrane domain. *Biochem Biophys Res Commun* 187:1135-1143
- ilotti 314. Graves PN, Magnusson RP, Fong P, Davies TF, Identification of a TSH variant protein lacking the transmembrane region in human serum. Program of the 75th Annual Meeting of The Endocrine Society, Las Vegas, NV, 1993 (Abstract 355)
- oglio 315. De Groot LJ, Sridama V 1989 Thyroid neoplasia. In: De Groot LJ, Bessen GM, Cahill Jr GF, Marshall JC, Nelson DH, Odell WD, Potts Jr JT (eds) *Endocrinology*. WB Saunders, Philadelphia, vol 2:758-776
- ne in 316. De Micco C, Ruf J, Chrestin MA, Gros N, Henry JF, Carayon P 1991 Immunohistochemical study of thyroid peroxidase in normal, hyperplastic and neoplastic thyroid tissue. *Cancer* 67:3036-3041
- Tissue 317. Monden T, Yamada M, Satoh T, Ilzuka M, Mori M (1992) Analysis of the TSH receptor gene structure in various thyroid disorders: DNA from thyroid can have large insertions or deletions. *Thyroid* 2:189-192
- ptide 318. Ling L, Haraguchi K, Ohta K, Enolo T, Onaya T 1992 β_2 adrenergic receptor mRNA is over-expressed in neoplastic human thyroid tissue. *Endocrinology* 130:547-549
- 1252- 319. Allen LF, Lefkowitz RJ, Caron MG, Cotecchia S 1991 G-protein-coupled genes as proto-oncogenes: constitutively activating mutation of the α_1 adrenergic receptor enhances mutagenesis and tumorigenicity. *Proc Natl Acad Sci USA* 86:11354-8
- uper- 319a. Parma J, Duprez L, Van Sande J, Cochaux P, Gervy C, Mockel J, Dumont J, Vassart G 1993 Somatic mutations in the thyrotropin receptor gene cause hyperfunctioning thyroid adenomas. *Nature* 365:649-651
- ature 319b. Lefkowitz RJ 1993 G protein coupled receptors: turned on to ill effect. *Nature* 365:603-604
320. Harris H 1988 The analysis of malignancy by cell fusion: the position in 1988. *Cancer Res* 48:3302-3306
321. Sager R 1985 Genetic suppression of tumor formation. *Adv Cancer Res* 44:43-68
322. Benedict WJ, Weissman BE, Mark C, Stanbridge EJ 1984 Tumorigenicity of human HT1080 fibrosarcoma X normal fibroblast hybrids: chromosome dosage dependency. *Cancer Res* 44:3471-3479
323. Harris H, Miller OJ, Klein G, Worst P, Tachibana T 1969 Suppression of malignancy by cell fusion. *Nature* 223:363-368
324. Knudson AG 1971 Mutation and cancer: statistical study of retinoblastoma. *Proc Natl Acad Sci USA* 68:820-823
325. Cavenee WK, Murphree AL, Shull MM, Benedict WF, Parkes RS, Kock E, Nordenskjold M 1986 Prediction of familial predisposition to retinoblastoma. *N Engl J Med* 314:1201-1207
326. Cavenee WK, Dryja TP, Phillips RA, Benedict WF, Godbout R, Gallie BL, Murphree AL, Strong LC, White RL 1983 Expression of recessive alleles by chromosomal mechanisms in retinoblastoma. *Nature* 305:779-784
327. Saxon PJ, Srivatsan GS, Stanbridge EJ 1986 Introduction of human chromosome 11 via microcell transfer controls tumorigenic expression of HeLa cells. *EMBO J* 5:3461-3466
328. Weissman BE, Saxon PJ, Pasquale SR, Jones GR, Geiser AG, Stanbridge EJ 1987 Introduction of normal human chromosome 11 into a Wilms' tumor cell line controls its tumorigenic expression. *Science* 236:175-180
329. Trent J, Stanbridge EJ, McBride HL, Meese GU, Casey G, Araujo DE, Witkowski CM, Neagle RB 1990 Tumorigenicity in human melanoma cell lines controlled by introduction of human chromosome 6. *Science* 247:568-571
330. Tanaka K, Oshimura M, Kikuchi R, Seki M, Hayashi T, Miyaki M 1991 Suppression of tumorigenicity in human colon carcinoma cells by introduction of normal chromosome 5 or 18. *Nature* 349:340-342
331. Shimizu M, Yokota J, Mori N, Shuin T, Shinoda M, Torada M, Oshimura M 1990 Introduction of normal chromosome 3p modulates the tumorigenicity of a human renal cell carcinoma cell line YC R. *Oncogene* 5:185-194
332. Dowdy SF, Fasching CL, Araujo D, Lai K-M, Livanos E, Weissman BE, Stanbridge EJ 1991 Suppression of tumorigenicity in Wilms tumor by the p15.5-p14 region of chromosome 11. *Science* 254:293-295
333. Bookstein R, Shew JY, Chen PL, Scully P, Lee WH 1990 Suppression of tumorigenicity of human prostate carcinoma cells by replacing a mutated RB gene. *Science* 247:712-715
334. Ponder B 1988 Cancer gene losses in human tumours [news]. *Nature* 335:400-402
335. Killary AM, Wolf ME, Giambernardi TA, Naylor SL 1992 Definition of a tumor suppressor locus within human chromosome 3p21-22. *Proc Natl Acad Sci USA* 89:9705-9709
336. Weiland I, Bohm M, Bogtaz S 1992 Isolation of DNA sequences deleted in lung cancer by genomic difference cloning. *Proc Natl Acad Sci USA* 89:9705-9709
337. Liang P, Pardee AB 1992 Differential display of eukaryotic messenger RNA by means of the polymerase chain reaction. *Science* 257:967-971
338. Herrmann MA, Hay ID, Bartelt Jr DH, Ritland SR, Dahl RJ, Grant CS, Jenkins RB 1991 Cytogenetic and molecular genetic studies of follicular and papillary thyroid cancers. *J Clin Invest* 88:1596-1604
339. Naylor SL, Johnson BE, Minna JD, Sakaguchi AY 1987 Loss of heterozygosity of chromosome 3p markers in small cell lung cancer. *Nature* 329:451-454
340. Kok K, Osinga J, Carritt B, Davis MB, van der Hout AH, van der Veen AY, Landsvater RM, de Leij LF, Berendsen HH, Postmus PE, Poppema S, Buys CHCM 1987 Deletion of a DNA sequence at the chromosomal region 3p21 in all major types of lung cancer. *Nature* 330:578-581
341. Baruch H, Johnson B, Hovis J, Yano T, Gazdar A, Pettengill OS, Graziano S, Sorenson GD, Poiesz BJ, Minna J, Linehan M, Zbar B 1987 The molecular analysis of the short arm of chromosome 3

- in small-cell and non-small cell carcinoma of the lung. *N Engl J Med* 317:1109-1113
342. Devilee P, van den Broek M, Kuipers-Dijkshoorn N, Kolluri R, Khan PM, Pearson PL, Cornelisse CJ 1989 At least four different chromosomal regions are involved in loss of heterozygosity in human breast carcinoma. *Genomics* 5:554-560
 343. Ali IU, Lidereau R, Callahan R 1989 Presence of two members of *c-erbA* receptor gene family (*c-erbA* beta and *c-erbA2*) in smallest region of somatic homozygosity on chromosome 3p21-p25 in human breast carcinoma. *J Natl Cancer Inst* 81:1571-1575
 344. Lothe RA, Fossa SD, Stenwig AE, Nakamura Y, White R, Børresen AL, Brogger A 1989 Loss of 3p or 11p alleles is associated with testicular cancer tumors. *Genomics* 5:134-138
 345. Kovacs G, Erlandsson R, Boldog F, Ingvarsson S, Muller-Brechlin R, Klein G, Sumegi J 1988 Consistent chromosome 3p deletion and loss of heterozygosity in renal cell carcinoma. *Proc Natl Acad Sci USA* 85:1571:1575
 346. Matsuo K, Tang SH, Fagin JA 1991 Allelotype of human thyroid tumors: loss of chromosome 11q13 sequences in follicular neoplasms. *Mol Endocrinol* 5:1873-1879
 347. Larsson C, Skogseid B, Oberg K, Nakamura Y, Nordenskjöld M 1988 Multiple endocrine neoplasia type I maps to chromosome 11 and is lost in insulinoma. *Nature* 332:85-87
 348. Bystrom C, Larsson C, Blomberg C, Sandelin K, Falkmer U, Skogseid B, Oberg K, Werner S, Nordenskjöld M 1990 Localization of the *MEN1* gene to a small region within chromosome 11q13 by deletion mapping in tumors. *Proc Natl Acad Sci USA* 87:1968-1972
 349. Bale AE, Norton JA, Wong EL, Fryburg JS, Maton PN, Oldfield EH, Streeten E, Aurbach GD, Brandi ML, Friedman E, Spiegel AM, Taggart RT, Marx SJ 1991 Allelic loss on chromosome 11 in hereditary and sporadic tumors related to familial multiple endocrine neoplasia type I. *Cancer Res* 51:1154-1157
 350. Friend SH, Horowitz JM, Gerber MR, Wang X-F, Bogenmann E, Li FP, Weinberg RA 1987 Deletions of a DNA sequence in retinoblastoma and mesenchymal tumors: organizations of the sequence and its encoded protein. *Proc Natl Acad Sci USA* 84:9059-9063
 351. Hong FD, Huang H-JS, To H, Young L-JS, Oro A, Bookstein R, Lee EY-HP, Lee W-H 1989 Structure of the retinoblastoma gene. *Proc Natl Acad Sci USA* 86:5502-5506
 352. Lee WH, Shew JY, Hong FD, Sery TW, Donoso LA, Young LJ, Bookstein R, Lee EY 1989 The retinoblastoma susceptibility gene encodes a nuclear phosphoprotein associated with DNA binding activity. *Nature* 329:642-645
 353. Whyte P, Buchkovich KJ, Horowitz JM, Friend SH, Raybuck M, Weinberg RA, Harlow E 1988 Association between an oncogene and an anti-oncogene: the adenovirus E1A proteins bind to the retinoblastoma gene product. *Nature* 334:124-129
 354. Egan C, Bayley ST, Branton PE 1989 Binding of the Rb1 protein to E1A products is required for adenovirus transformation. *Oncogene* 4:383-388
 355. De Caprio JA, Ludlow JW, Figge J, Shew JY, Huang CM, Lee WH, Marsilio E, Paucha E, Livingston DM 1988 SV40 large tumor antigen forms a specific complex with the product of the retinoblastoma susceptibility gene. *Cell* 54:275-283
 356. Munger K, Werness BA, Dyson N, Phelps WC, Harlow E, Howley PM 1989 Complex formation of human papillomavirus E7 proteins with the retinoblastoma tumor suppressor gene product. *EMBO J* 8:4099-4105
 357. Figge J, Webster T, Smith TF, Paucha E 1988 Prediction of similar transforming regions in simian virus 40 large T, adenovirus E1A and myc oncoproteins. *J Virol* 62:1814-1818
 358. Hu Q, Dyson N, Harlow E 1990 The regions of the retinoblastoma protein needed for binding to adenovirus E1A or SV40 large T antigen are common sites for mutations. *EMBO J* 9:1147-1155
 359. Ludlow JW, DeCaprio JA, Huang CM, Lee WH, Paucha E, Livingston DM 1989 SV40 large T antigen binds preferentially to an underphosphorylated member of the retinoblastoma susceptibility gene product family. *Cell* 56:57-65
 360. De Caprio JA, Ludlow JW, Lynch D, Furukawa Y, Griffin J, Piwnicka-Worms H, Huang CM, Livingston DM 1989 The product of the retinoblastoma susceptibility gene has properties of a cell cycle regulatory element. *Cell* 58:1085-1095
 361. Mihara K, Cao XR, Yen A, Chandler S, Driscoll B, Murphree AL, T'Ang A, Fung YK 1989 Cell cycle-dependent regulation of phosphorylation of the human retinoblastoma gene product. *Science* 246:1300-1303
 362. Buchkovich K, Duffy LA, Harlow E 1989 The retinoblastoma protein is phosphorylated during specific phases of the cell cycle. *Cell* 58:1097-1105
 363. Stein GH, Beeson M, Gordon L 1990 Failure to phosphorylate the retinoblastoma gene product in senescent human fibroblasts. *Science* 249:666-669
 364. Furukawa Y, DeCaprio JA, Freedman A, Kanakura Y, Nakamura M, Ernst TJ, Livingston DM, Griffin JD 1990 Expression and state of phosphorylation of the retinoblastoma susceptibility gene product in cycling and noncycling human hematopoietic cells. *Proc Natl Acad Sci USA* 87:2770-2774
 365. Chellappan SP, Hiebert S, Mudryj M, Horowitz JM, Nevins JR 1991 The E2F transcription factor is a cellular target for the Rb protein. *Cell* 65:1053-1061
 366. Bagchi S, Weinmann R, Raychaudhuri P 1991 The retinoblastoma protein copurifies with E2F-L, an E1A-regulated inhibitor of the transcription factor E2F. *Cell* 65:1063-1072
 367. Chittenden T, Livingston DM, Kaelin Jr WG 1991 The T/E1A-binding domain of the retinoblastoma product can interact selectively with a sequence-specific DNA-binding protein. *Cell* 65:1073-1082
 368. Helin KA 1992 cDNA encoding a pRB-binding protein with properties of the transcription factor E2F. *Cell* 70:337-350
 369. Kaelin Jr WG, Krek W, Sellers WR, DeCaprio JA, Ajchenbaum F, Fuchs CS, Chittenden T, Li Y, Farnham PJ, Blarar MA, Livingston DM, Flemington EK 1992 Expression cloning of a cDNA encoding a retinoblastoma-binding protein with E2F-like properties. *Cell* 70:351-364
 370. Nevins JR 1992 E2F: a link between the Rb tumor suppressor protein and viral oncoproteins. *Science* 258:424-429
 371. Dowdy SF, Hinds PW, Louie K, Reed SI, Arnold A, Weinberg RA 1993 Physical interaction of the retinoblastoma protein with human D cyclins. *Cell* 73:499-511
 372. Ewen ME, Sluss HK, Sherr CJ, Matsushime H, Kato J-Y, Livingston DM 1993 Functional interactions of the retinoblastoma protein with mammalian D-type cyclins. *Cell* 73:487-497
 373. Devoto SH, Mudryj MM, Pines J, Hunter T, Nevins JR 1992 A cyclin A-protein kinase complex possesses sequence-specific DNA binding activity: p53^{cdk2} is a component of the E2F-cyclin A complex. *Cell* 68:167-176
 374. Cao L, Faha B, Dembski M, Tsai LH, Harlow E, Dyson N 1992 Independent binding of the retinoblastoma protein and p107 to the transcription factor E2F. *Nature* 355:176-179
 375. Shirodkar S, Ewen M, DeCaprio JA, Morgan J, Livingston DM, Chittenden T 1992 The transcription factor E2F interacts with the retinoblastoma product and a p107-cyclin A complex in a cell cycle-regulated manner. *Cell* 68:157-166
 376. Rustgi AK, Dyson N, Bernards R 1991 Amino-terminal domains of *c-myc* and *N-myc* proteins mediate binding to the retinoblastoma gene product. *Nature* 352:541-544
 377. Goodrich DW, Lee W-H 1992 Abrogation by *c-myc* of G1 phase arrest induced by Rb protein but not by p53. *Nature* 360:177-179
 378. Clarke AR, Maandag ER, van Roon M, van der Lugt NM, van der Valk M, Hooper ML, Berns A, te Riele H 1992 Requirement for a functional Rb-1 gene in murine development. *Nature* 359:328-330
 379. Lee EY-HP, Chang CY, Hu N, Wang YCJ, Lai CC, Herrup K, Lee WH, Bradley A 1992 Mice deficient for Rb are nonviable and show defects in neurogenesis and haematopoiesis. *Nature* 359:218-221
 380. Jacks T, Fazeli A, Schmitt EM, Bronson RT, Goodell MA, Weinberg RA 1992 Effects of an Rb mutation in the mouse. *Nature* 359:295-300
 381. Harlow E 1992 For your eyes only. *Nature* 359:270-271
 382. Lee EY-HP, Huang S, Shew JY, Donoso LA, Lee WH 1991 Diverse mutations lead to inactivation of the retinoblastoma gene. *Int Rev Cytol*

- ies of a
- arphree
ation of
ct. Sci-
- lastoma
il cycle.
- late the
sts. Sci-
- Naka-
ression
ptibility
nc cells.
- vins JR
the RB
- noblas-
bitor of
- T/E1A-
t selec-
a. Cell
- h prop-
- nbaum
ir MA,
ig of a
2F-like
- pressor
- inberg
human
- Y, Liv-
astoma
- 1992 A
c DNA
clin A
- √ 1992
107 to
- n DM,
ith the
a cell
- omains
astoma
- l phox
7-179
- A, van
ement
Nature
- K, Lee
e and
Nature
- Wein-
ature
- iverse
Faber
- DB, Chader GJ (eds) *Molecular Biology of the Retina: Basic and Clinically Relevant Studies*. Wiley-Liss Inc, New York, pp 221-240
383. Jensen RD, Miller RW 1971 Retinoblastoma: epidemiologic characteristics. *N Engl J Med* 285:307-311
 384. Hansen MF, Koufos A, Gallie BL, Phillips RA, Fodstad O, Brogger A, Gedde-Dahl T, Caveness WK 1985 Osteosarcoma and retinoblastoma: a shared chromosomal mechanism revealing recessive predisposition. *Proc Natl Acad Sci USA* 82:6216-6220
 385. Scheffner M, Munger K, Byrne JC, Howley PM 1991 The state of the p53 and retinoblastoma genes in human cervical carcinoma cell lines. *Proc Natl Acad Sci USA* 88:5523-5527
 386. Rygaard K, Sorenson GD, Pettengill OS, Cate CC, Spang-Thomsen M 1990 Abnormalities in structure and expression of the retinoblastoma gene in small cell lung cancer cell lines and xenografts in nude mice. *Cancer Res* 50:5312-5317
 387. Cheng J, Scully P, Shew JY, Lee WH, Vila V, Haas M 1990 Homozygous deletion of the retinoblastoma gene in an acute lymphoblastic leukemia (T) cell line. *Blood* 75:730-735
 388. Harbour JW, Lai SL, Whang-Peng J, Gazdar AF, Minna JD, Kaye FJ 1988 Abnormalities in structure and expression of the human retinoblastoma gene in SCLC. *Science* 21:353-357
 389. Horowitz JM, Yandell DW, Park SH, Canning S, Whyte P, Buchkovich K, Harlow E, Weinberg RA, Dryja TP 1989 Point mutational inactivation of the retinoblastoma antioncogene. *Science* 243:937-940
 390. Horowitz JM, Park SH, Bogenmann E, Cheng JC, Yandell DW, Kaye FJ, Minna JD, Dryja TP, Weinberg RA 1990 Frequent inactivation of the retinoblastoma anti-oncogene is restricted to a subset of human tumor cells. *Proc Natl Acad Sci USA* 87:2775-2779
 391. Lee EY, To H, Shew JY, Bookstein R, Scully P, Lee WH 1988 Inactivation of the retinoblastoma susceptibility gene in human breast cancers. *Science* 241:218-221
 392. Tang A, Wu KJ, Hashimoto T, Liu W-Y, Takahashi R, Shi X-H, Mihara K, Zhang F-H, Chen YY, Du C, Qian J, Lin Y-G, Murphree AL, Qiu W-R, Thompson T, Benedict WF, Fung Y-KT 1989 Genomic organization of the human retinoblastoma gene. *Oncogene* 4:401-407
 393. Xu H-J, Hu SX, Hashimoto T, Takahashi R, Benedict WF 1989 The retinoblastoma susceptibility gene product: a characteristic pattern in normal cells and abnormal expression in malignant cells. *Oncogene* 4:807-812
 394. Yokota J, Akiyama T, Fung YKT, Benedict WF, Namba Y, Hanaoka M, Wada M, Terasaki T, Shimozato Y, Sugimura T, Terada M 1988 Altered expression of the retinoblastoma (RB) gene in small-cell carcinoma of the lung. *Oncogene* 4:471-475
 395. Reisman P, Simon MA, Lee W-H, Salmon DJ 1989 Studies of the retinoblastoma gene in human sarcoma. *Oncogene* 4:839-843
 396. Toguchida J, Ishizaki K, Sasaki MS, Ikenaga M, Sugimoto M, Kotoura Y, Yamamuro T 1988 Chromosomal reorganization for the expression of recessive mutation of retinoblastoma susceptibility gene in the development of osteosarcoma. *Cancer Res* 48:3939-3943
 397. Hu QJ, Dyson N, Harlow E 1990 The regions of the retinoblastoma protein needed for binding to adenovirus E1A or SV40 large T antigen are common sites for mutations. *EMBO J* 9:1147-1155
 398. Huang S, Wang N-P, Tseng BY, Lee W-H, Lee EHH-P 1990 Two distinct and frequently mutated regions of retinoblastoma protein are required for binding of SV40 T antigen. *EMBO J* 9:1815-1822
 399. Kaelin WG, Ewen ME, Livingston DM 1990 Definition of the minimal simian virus 40 large T antigen and adenovirus E1A-binding domain in the retinoblastoma gene product. *Mol Cell Biol* 10:3761-3769
 400. Dunn JM, Phillips A, Becker AJ, Gallie BL 1988 Identification of germline and somatic mutations affecting the retinoblastoma gene. *Science* 241:1797-1800
 401. Shew JY, Lin BTY, Chen PL, Tseng BY, Yang-Feng TL, Lee WH 1990 C-terminal truncation of the retinoblastoma gene product leads to functional inactivation. *Proc Natl Acad Sci USA* 87:6-10
 402. Ledent C, Dumont J, Vassart G, Parmentier M 1991 Thyroid adenocarcinomas secondary to tissue-specific expression of SV40 large T antigen in SV40-transformed cells. *Nature* 278:261-263
 403. Zou MJ, Shi YF, Farid NR 1993 Frequent inactivation of the retinoblastoma gene in human thyroid carcinoma. *Endocrinol*, in press
 404. Soussi T, Carom de Formentel C, May P 1990 Structural aspects of the p53 protein in relation to gene evolution. *Oncogene* 5:945-952
 405. Kern SE, Kinzler KW, Bruskin A, Jarosz D, Friedman P, Prives C, Vogelstein B 1991 Identification of p53 as a sequence-specific DNA-binding protein. *Science* 252:1708-1711
 406. Milner J, Medcalf EA 1991 Cotranslation of activated mutant p53 with wild type drives the wild-type p53 protein into the mutant conformation. *Cell* 65:765-774
 407. Stenger JE, Mayr GA, Mann K, Tegtmeyer P 1992 Formation of stable p53 homotetramers and multiples of tetramers. *Mol Carcinog* 5:102-106
 408. Kern SE, Pietenpol JA, Thiagalingam S, Seymour A, Kinzler KW, Vogelstein B 1992 Oncogenic forms of p53 inhibit p53-regulated gene expression. *Science* 256:827-830
 409. Unger T, Nau MM, Segal S, Minna JD 1992 p53: a transdominant regulator of transcription whose function is ablated by mutations occurring in human cancer. *EMBO J* 11:1383-1390
 410. Miller CW, Imai Y, Aslo A, Li L, Koeffler HP 1992 Sublocalization of transcriptional activation domain of p53. *Proc Am Assoc Cancer Res* 33:386
 411. Raycroft L, Wu HY, Lozano G 1990 Transcriptional activation by wild type but not transforming mutants of the p53 antioncogene. *Science* 249:1049-1051
 412. Fields S, Jang SK 1990 Presence of a potent transcriptional activation sequence in the p53 protein. *Science* 249:1046-1049
 413. Hinds PW, Finlay CA, Frey AB, Levine AJ 1987 Immunological evidence for the association of p53 with a heat shock protein, hsc70, in p53-plus-ras-transformed cell lines. *Mol Cell Biol* 7:2863-2869
 414. Shaulsky G, Goldfinger N, Ben Ze'ev A, Rotter V 1990 Nuclear accumulation of p53 protein is mediated by several nuclear localization signals and plays a role in tumorigenesis. *Mol Cell Biol* 10:6565-6577
 415. Bischoff JR, Friedman PN, Marshak DR, Prives C, Beach D 1990 Human p53 is phosphorylated by p60-cdc2 and cyclin B-cdc2. *Proc Natl Acad Sci USA* 87:4766-4770
 416. Zou M, Shi Y, Farid NR 1993 p53 mutations in all stages of thyroid carcinomas. *J Clin Endocrinol Metab* 77:1054-1058
 417. Kern SE, Kinzler KW, Baker SJ, Nigro JM, Rotter V, Levine AJ, Friedman P, Prives C, Vogelstein B 1991 Mutant p53 proteins bind DNA abnormally *in vitro*. *Oncogene* 6:131-136
 418. Milner J, Watson JV 1990 Addition of fresh medium induces cell cycle land conformation changes in p53, a tumour suppressor protein. *Oncogene* 5:1683-1690
 419. Weintraub B, Hauschka S, Tapscott SJ 1991 The MCK enhancer contains a p53 responsive element. *Proc Natl Acad Sci USA* 88:4570-4571
 420. Agoff SN, Hov J, Linzer DH, Wu B 1993 Regulation of the human hsp70 promoter by p53. *Science* 259:84-87
 421. Weinert TA, Hartwell LH 1988 The RAD9 gene controls the cell cycle response to DNA damage in *Saccharomyces cerevisiae*. *Science* 241:317-322
 422. Malkin D, Li DFP, Strong LC, Fraumeni JF, Nelson CE, Kim DH, Kassel J, Gryka MA, Bischoff FZ, Tainsky MA, Friend SH 1990 Germline mutation in a familial syndrome of breast cancer, sarcomas and other neoplasms. *Science* 250:1233-1238
 423. Maltzman W, Czyzk L 1984 UV irradiation stimulates levels of p53 cellular tumor antigen in non-transformed mouse cells. *Mol Cell Biol* 4:1689-1694
 424. Lu X, Park SH, Thompson TC, Lane DP 1992 ras-induced hyperplasia occurs with mutation of p53, but activated ras and myc together can induce carcinoma without p53 mutations. *Cell* 70:153-161
 425. Linzer DL, Levine AJ 1979 Characterization of a 54 K dalton cellular SV40 tumor antigen present in SV40 transformed cells and in uninfected embryonal cells. *Cell* 17:43-52
 426. Eliyahu D, Raz A, Gruss P, Givol D, Oren M 1984 Participation

- of p53 cellular tumour antigen in transformation of normal embryonic cells. *Nature* 312:646-649
427. Parada LF, Land H, Weinberg RA, Wolf D, Rotter V 1984 Cooperation between gene encoding p53 tumour antigen and ras in cellular transformation. *Nature* 312:649-651
 428. Levine AJ, Momand J, Finlay CA 1991 The p53 tumor suppressor gene. *Nature* 351:453-455
 429. Finlay CA, Hinds PW, Tan TH, Elihayu D, Oren M, Levine AJ 1988 Activating mutations for transformation by p53 produce a gene product that forms a hsc70-p53 complex with an altered half-life. *Mol Cell Biol* 8:531-539
 430. Lane DP, Crawford LV 1979 T-antigen is bound to host protein in SV40-transformed cells. *Nature* 278:261-263
 431. Jenkins JR, Chumakov P, Addison C, Sturzbecher HW, Wade-Evans A 1988 Two distinct regions of the murine p53 primary amino acid sequence are implicated in stable complex formation with simian virus 40 T antigen. *J Virol* 62:3903-3906
 432. Sarnow P, Ho YS, Williams J, Levine AJ 1982 Adenovirus E1b-58kd tumor antigen and SV40 large tumor antigen are physically associated with the same 54 kd cellular protein in transformed cells. *Cell* 28:387-394
 433. Werness BA, Levine AJ, Howley PM 1990 Association of human papillomavirus types 16 and 18 E6 proteins with p53. *Science* 248:76-79
 434. Li FP, Fraumeni Jr JF, Mulvihill JJ, Blattner WA, Dreyfus MG, Tucker MA, Miller RW 1988 A cancer family syndrome in twenty-four kindreds. *Cancer Res* 48:5348-5362
 435. Garber JE, Goldstein AM, Kantor AF, Fraumeni JF, Li FP 1991 Follow up study of twenty four families with Li Fraumeni syndrome. *Cancer Res* 51:6094-6097
 436. Malkin D, Jolly KW, Barbier N, Look AT, Friend SH, Gebhardt MC, Andersen TI, Borresen AL, Li FP, Garber J, Greenberg ML, Strong LC 1992 Germline mutations of the p53 tumor-suppressor gene in children and young adults with second malignant neoplasms. *N Engl J Med* 326:1309-1315
 437. Toguchida J, Yamaguchi T, Dayton SH, Beauchamp RL, Herrera GE, Ishizaki K, Yamamoto T, Meyers PA, Little JB, Sasaki MS, Weichselbaum RR, Yandell DW 1992 Prevalence and spectrum of germline mutations of the p53 gene among patients with sarcomas. *N Engl J Med* 326:1301-1308
 438. Iavarone A, Matthay KK, Steinkrchner TM, Israel MA 1992 Germ-line and somatic p53 gene mutations in multifocal osteogenic sarcoma. *Proc Natl Acad Sci USA* 89:4207-4209
 439. Metzger AK, Sheffield VC, Duyk G, Daneshvar L, Edwards MSB, Cogen PH 1991 Germ-line mutation in the p53 gene in a patient with an intracranial ependymoma. *Proc Natl Acad Sci USA* 88:7825-7829
 440. Sidransky D, Tokino T, Helzlsouer K, Zehnbauser B, Raush G, Shelton B, Prestigiacomo L, Vogelstein B, Davidson N 1992 Inherited p53 gene mutations in breast cancer. *Cancer Res* 52:2984-2986
 441. Borresen AL, Andersen TI, Garber J, Barbier N, Thorlacius S, Eyfjord J, Ottestad L, Sorensen BS, Pharm M, Hovig G, Friend SH 1992 Screening for germ-line p53 mutations in breast cancer patients. *Cancer Res* 52:3234-3236
 442. Srivastava S, Zou Z, Pirolo K, Blattner W, Change EH 1990 Germline transmission of a mutated p53 gene in a cancer-prone family with Li-Fraumeni syndrome. *Nature* 348:747-749
 443. Frebourg T, Friend SH 1992 Cancer risks from germline p53 mutations. *J Clin Invest* 90:1637-1641
 444. Hollstein M, Sidransky D, Vogelstein B, Harris CC 1991 p53 mutations in human cancers. *Science* 253:49-53
 445. Nigro JM, Baker SJ, Preisinger AC, Jessup JM, Hostetter R, Cleary K, Bigner SH, Davidson N, Baylin S, Devilee P, Glover T, Collins FS, Weston A, Modali R, Harris CC, Vogelstein B 1989 Mutations in the p53 gene occur in diverse human tumour types. *Nature* 342:705-708
 446. Sidransky D, Mikkelsen T, Schwechheimer K, Rosenblum ML, Cavenee W 1992 Clonal expansion of p53 mutant cells is associated with brain tumor progression. *Nature* 355:846-847
 447. Davidoff AM, Kerns BJM, Iglehart JD, Marks JR 1991 Maintenance of p53 alterations throughout breast cancer progression. *Cancer Res* 51:2605-2610
 448. Takahashi T, Nau NM, Chiba I, Birrer MJ, Rosenberg RK, Vinocour M, Levitt M, Pass H, Gazdar AF, Minna JD 1989 p53 - A frequent target for genetic abnormalities in lung cancer. *Science* 246:491-494
 449. Hollstein MC, Metcalf RA, Welsh JA, Montesano R, Harris CC 1990 Frequent mutation of the p53 gene in human esophageal cancer. *Proc Natl Acad Sci USA* 87:9958-9961
 450. Sidransky D, Voneschenbach A, Tsai YC, Jones P, Summerhayes I, Marshall F, Paul M, Green P, Hamilton SR, Frost P, Vogelstein B 1991 Identification of p53 gene mutations in bladder cancers and urine samples. *Science* 252:706-709
 451. Bartek J, Bartkova J, Vojtesek B, Staskova Z, Lukas J, Rejthar A, Kovarik J, Midgley CA, Gannon JV, Lane DP 1991 Aberrant expression of the p53 oncoprotein is a common feature of a wide spectrum of human malignancies. *Oncogene* 6:1699-1703
 452. Frebourg T, Kassel J, Lam KT, Gryka MA, Barbier N, Andersen TI, Borresen AL, Friend SH 1992 Germline mutations of the p53 tumor suppressor gene in patients with high risk for cancer inactivate the p53 protein. *Proc Natl Acad Sci USA* 89:6413-6417
 453. Carbone D, Chiba I, Mitsudomi T 1991 Polymorphism at codon 213 within the p53 gene. *Oncogene* 6:1691-1692
 454. Matlashewski GJ, Tuck S, Pim D, Lamb P, Schneider J, Crawford LV 1987 Primary structure polymorphism at amino acid residue 72 of human p53. *Mol Cell Biol* 7:961-963
 455. Hinds PW, Finlay CA, Quartin RS, Baker SJ, Fearn ER, Vogelstein B, Lemine AJC 1990 Mutant p53 DNA clones for human colon carcinomas cooperate with ras in transforming primary rat cells: a comparison of the 'hot spot' mutant phenotypes. *Cell Growth Differentiation* 1:571-580
 456. Stephen CW, Lane DP 1992 Mutant conformation of p53. Precise epitope mapping using a filamentous phage epitope library. *J Mol Biol* 225:577-583
 457. Hsu IC, Metcalf RA, Sun T, Welsh JA, Wang NJ, Harris CC 1991 Mutational hotspot in the p53 gene in human hepatocellular carcinoma. *Nature* 350:427-428
 458. Vogelstein B, Kinzler KW 1992 Carcinogens leave fingerprints. *Nature* 355:209-210
 459. Brash DE, Rudolph JA, Simon JA, Lin A, McKenna GJ, Baden HP, Halpern AJ, Poter J 1991 A role for sunlight in skin cancer: UV-induced p53 mutations in squamous cell carcinoma. *Proc Natl Acad Sci USA* 88:10124-10128
 460. Momand J, Zambetti GP, Olson DC, George DL, Levine AJ 1992 The mdm-2 oncogene product forms a complex with the p53 protein and inhibits p53-mediated transactivation. *Cell* 69:1237-1245
 461. Oliner JD, Kinzler KW, Meltzer PS, George DL, Vogelstein B 1992 Amplification of a gene encoding a p53-associated protein in human sarcomas. *Nature* 358:80-83
 462. Baker SJ, Preisinger AC, Jessup JM, Paraskeva C, Markowitz S, Wilson JK, Hamilton S, Vogelstein B 1990 p53 gene mutations occur in combination with 17p allelic deletions as late events in colorectal tumorigenesis. *Cancer Res* 50:7717-7722
 463. Bennett WP, Hollstein MC, He A, Zhu SM, Resau JH, Trump BF, Metcalf RA, Welsh JA, Midgley C, Lane DP, Harris CC 1991 Archival analysis of p53 genetic and protein alteration in Chinese esophageal cancer. *Oncogene* 6:1779-1784
 464. Watling DL, Gown AM, Coltrera MD 1992 Overexpression of p53 in head and neck cancer. *Head Neck* 14:437-444
 465. Wright PA, Lemoine NR, Goretzki PE, Wyllie FS, Bond J, Hughes C, Roher HD, Williams ED, Wynford-Thomas D 1991 Mutation of the p53 gene in a differentiated human thyroid carcinoma cell line, but not in primary thyroid tumours. *Oncogene* 6:1693-1697
 466. Fagin JA, Matuso K, Karmakar A, Lin Chew D, Tang S-H, Koeffler HP 1992 High prevalence of mutations of the p53 gene in poorly differentiated human carcinoma. *J Clin Invest* 91:179-184
 467. Ito T, Seyama T, Mizuno T, Tsuyama N, Hayashi T, Hayashi Y, Dohi K, Nakamura N, Akiyama M 1992 Unique association of

- p53 mutations with undifferentiated but not with differentiated carcinomas of the thyroid gland. *Cancer Res* 52:1369-1371
468. Rideout III WM, Coetzee GA, Olumi AF, Jones PA 1990 5-Methylcytosine as an endogenous mutagen in the human LDL receptor and p53 genes. *Science* 249:1288-1290
 469. Shen J-C, Rideout III WM, Jones PA 1992 High frequency mutagenesis by a DNA methyl transferase. *Cell* 71:1073-1080
 470. Ahuja HG, Testa MP, Cline MJ 1990 Variation in the protein coding region of the human p53 gene. *Oncogene* 5:1409-1410
 471. Harris N, Brill E, Shohat O, Prokocimer M, Wolf D, Arai N, Rotter V 1986 Molecular basis for heterogeneity of the human p53 protein. *Mol Cell Biol* 6:4650-4656
 472. Kerbel RS 1992 Expression of multi-cytokine resistance and multi-growth factor independence in advanced stage metastatic cancer. *Am J Pathol* 141:519-524
 473. Steeg PS, Bevilacqua G, Kopper L, Thorgerisson UP, Talmadge JE, Liotta LA, Sobel ME 1988 Evidence for a novel gene associated with low tumor metastatic potential. *J Natl Cancer Inst* 80:200-204
 474. Kerbel RS, Waghorne C, Korczak B, Lagarde A, Breitman M 1988 Clonal dominance of primary tumors by metastatic cells: genetic analysis and biological implications. *Cancer Surv* 7:597-629
 475. Kerbel RS 1990 Growth dominance of the metastatic cancer cell: cellular and molecular aspects. *Adv Cancer Res* 55:87-132
 476. Halaban R, Kwon B, Ghosh S, Delli Bovis P, Baird A 1988 bFGF as an autocrine growth factor for human melanocytes. *Oncogene* Res 3:177-186
 477. Becker D, Meier CB, Herlyn M 1989 Proliferation of human malignant melanomas is inhibited by antisense oligodeoxynucleotides targeted against basic fibroblast growth factor. *EMBO J* 8:3685-3691
 478. Rodeck U, Becker D, Herlyn M 1991 Basic fibroblast growth factor in human melanomas. *Cancer Cells* 3:308-311
 479. Sporn MB, Todaro GJ 1980 Autocrine secretion and malignant transformation of cells. *N Engl J Med* 303:878-880
 480. Browder TM, Dunbar CE, Nienhuis AW 1989 Private and public autocrine loops in neoplastic cells. *Cancer Cells* 1:9-17
 481. Herlyn M 1990 Human melanoma: development and progression. *Cancer Metastasis Rev* 9:101-112
 482. Richmond A 1991 The pathogenic role of growth factors in melanoma. *Semin Dermatol* 10:246-255
 483. Schwarz LC, Gingras MC, Goldberg G, Breenberg AH, Wright JA 1988 Loss of growth factor dependence and conversion of transforming growth factor 1 inhibition to stimulation in metastatic H-ras transformed murine fibroblast. *Cancer Res* 48:6999-7003
 484. Mooradian DL, Purchio AF, Furcht LT 1990 Differential effects of transforming growth factor on the growth of poorly and highly metastatic murine melanoma cells. *Cancer Res* 50:273-277
 485. Theodorescu D, Caltabiano M, Greig R, Rieman D, Kerbel RS 1991 Reduction of TGF-beta activity abrogates growth promoting tumor cell-cell interactions *in vivo*. *J Cell Physiol* 148:380-390
 486. Watson JM, Sensintaffar JL, Berek JS, Martinez-Maza O 1990 Constitutive production of interleukin 6 by ovarian cancer cell lines and by primary ovarian tumor cultures. *Cancer Res* 50:6959-6965
 487. Takenawa J, Kaneko Y, Fukumoto M, Hirano T, Fukuyama H, Nakayama H, Fujita J, Yoshida O 1991 Enhanced expression of interleukin-6 in primary human renal cell carcinomas. *J Natl Cancer Inst* 83:1668-1672
 488. Siegall CB, Schwab G, Nordan RP, FitzGerald DJ, Pastan I 1990 Expression of the interleukin 6 receptor and interleukin 6 in prostate carcinoma cells. *Cancer Res* 50:7786-7788
 489. Van Meir E, Sawamura Y, Diserens A-C, Hamour M-F, de Tribolet N 1990 Human glioblastoma cells release interleukin 6 *in vivo* and *in vitro*. *Cancer Res* 50:6683-6688
 490. Tabibzadeh SS, Poubouridia D, May LT, Sehgal PB 1989 Interleukin-6 immunoreactivity in human tumors. *Am J Pathol* 135:427-433
 491. Miki S, Isawano M, Miki Y, Yamamoto M, Tang B, Yokokawa K, Sonoda T, Hirano T, Kishimoto T 1989 Interleukin-6(IL-6) functions as an *in vitro* autocrine growth factor in renal cell carcinomas. *FEBS Lett* 250:607-610
 492. Folkman J 1990 What is the evidence that tumors are angiogenesis dependent? *J Natl Cancer Inst* 82:4-6
 493. Bickell R, Harris AL 1991 Novel growth regulatory factors and tumor angiogenesis. *Eur J Cancer* 27:781-784
 494. Srivastava A, Laider P, Davies R, Horgan K, Hughes LE 1988 The prognostic significance of tumor vascularity in intermediate-thickness (0.76-4 mm thick) skin melanoma. *Am J Pathol* 133:419-423
 495. Horak GR, Leek R, Klenk N, Lejeune S, Smith K, Stuart N, Greenall M, Stepniowska K, Harris AL 1992 Angiogenesis, assessed by platelet/endothelial cell adhesion molecule antibodies as indicator of node metastases and survival in breast cancer. *Lancet* 340:1120-1124
 496. Steeg PS, Bevilacqua G, Rosengard AM, Sobel ME, Cioce V, Liotta LA 1989 Altered gene expression in tumor metastasis: the nm23 gene. In: Schirmacher V, Schwartz-Albiez R (eds) *Cancer Metastasis. Molecular and Cellular Biology, Host Immune Responses and Perspectives for Treatment*. Springer Verlag, Berlin, pp 48-52
 497. Steeg PS, Bevilacqua G, Pozzatti R, Liotta LA, Sobel ME 1988 Altered expression of nm23, a gene associated with low tumor metastatic potential, during adenovirus 2 Ela inhibition of experimental metastasis. *Cancer Res* 48:6550-6554
 498. McAlpine PJ, Shows TB, Boucheix C, Huebner M, Anderson WA 1991 The 1991 catalog of mapped genes and report of the nomenclature committee. *Cytogenet Cell Genet* 58:5-102
 499. Rosengard AM, Krutzsch HC, Shearn A, Biggs JR, Barker E, Margulies IMK, King CR, Liotta LA, Steeg PS 1989 Reduced Nm23/Awd protein in tumor metastasis and aberrant *Drosophila* development. *Nature* 342:177-180
 500. Wallet V, Mutzel R, Troll H, Barzu O, Wurster B, Veron M, Lacombe M-L 1990 Dictyostelium nucleotide diphosphate kinase highly homologous to nm23 and awd proteins involved in mammalian tumor metastasis and *Drosophila* development. *J Natl Cancer Inst* 82:1199-1202
 501. Kimura N, Shimada N, Nomura K, Watanabe K 1990 Isolation and characterization of a cDNA clone encoding rat nucleoside diphosphate kinase. *J Biol Chem* 265:15744-15749
 502. Biggs J, Hersperger E, Steeg PS, Liotta LA, Shearn A 1990 A *Drosophila* gene that is homologous to a mammalian gene associated with tumor metastasis codes for a nucleoside diphosphate kinase. *Cell* 63:933-940
 503. Stahl JA, Leona A, Rosengard AM, Porter L, King CR, Steeg PS 1991 Identification of a second human nm23 gene, nm23-H2. *Cancer Res* 51:445-449
 504. Gillet A-M, Presecq E, Vonica A, Lasco I 1991 Nucleoside diphosphate kinase from human erythrocytes. *J Biol Chem* 266:8784-8789
 505. Backer JM, Mendola CE, Kovacs I, Fairhurst JL, O'Hara B, Eddy Jr RL, Shows TB, Mathew S, Murty VVVS, Chaganti RSK 1993 Chromosomal localization and nucleoside diphosphate kinase activity of human metastasis-suppressor genes nm23-1 and nm23-2. *Oncogene* 8:497-502
 506. Bevilacqua G, Sobel ME, Liotta LA, Steeg PS 1989 Association of low nm23 RNA levels in human primary infiltrating ductal breast carcinomas with lymph node involvement and other histopathological indicators of high metastatic potential. *Cancer Res* 49:5185-5190
 507. Barnes R, Masood S, Barker E, Rosengard AM, Coggin DL, Crowell T, King CR, Porter-Jordan K, Wargotz ES, Liotta LA, Steeg PS 1991 Low nm23 protein expression in infiltrating ductal breast carcinomas correlates with reduced patient survival. *Am J Pathol* 139:245-250
 508. Beahrs OH, Henson DE, Hutter RVP 1988 Head and neck tumors. *Thyroid Gland Manual for Staging of Cancer*. American Joint Commission on Cancer
 509. Leone A, Flatow U, King CR, Sandeen MA, Margulies IMK, Liotta LA, Steeg PS 1991 Reduced tumor incidence, metastatic potential and cytokine responsiveness of nm23 transfected melanoma cells. *Cell* 65:25-35

510. Leone A, McBride OW, Weston A, Wang MG, Anglard P, Cropp CS, Goepel JR, Lidereau R, Callahan R, Linehan WM, Rees RC, Harris CC, Liotta LA, Steeg PS 1991 Somatic allelic deletion of nm23 in human cancer. *Cancer Res* 51:2490-2493
511. Cohn KH, Wang F, Desoto-Lapaix F, Solomon WB, Patterson LG, Arnold MR, Weimar J, Feldman JG, Levy AT, Leone A, Steeg PS 1991 Association of nm23-H1 allelic deletions with distant metastases in colorectal carcinoma. *Lancet* 338:722-724
512. Wang L, Patel V, Ghosh L, Chen H-C, Banarjee S 1993 Mutations in nm23 gene is associated with metastasis in colorectal cancer. *Cancer Res* 53:717-720
513. Haut M, Steeg PS, Willson JKV, Markowitz SD 1991 Induction of nm23 gene expression in human colonic neoplasms and equal expression in colon tumors of high and low metastatic potential. *J Natl Cancer Inst* 83:712-716
514. Hailat N, Keim DR, Melhem RF, Zhu X-X, Eckerskorn C, Brodeur GM, Reynolds CP, Seeger RC, Lottspeich F, Strahler JR, Hanash SM 1991 High levels of p19/nm23 protein in neuroblastoma are associated with advanced stage disease and with N-myc gene amplification. *J Clin Invest* 88:341-345
515. Lacombe M-L, Sastre-Garau X, Lascu I, Vonica A, Wallet V, Thiery JP, Veron M 1991 Overexpression of nucleoside diphosphate kinase (nm23) in solid tumors. *Eur J Cancer* 27:1302-1307
516. Zou MJ, Shi YF, Farid NR 1993 High levels of Nm23 gene expression in the advanced stage of thyroid cancer. *Br J Cancer* 68:385-388
517. Kikkawa S, Takahashi K, Shimada N, Vi M, Kimura N, Katada T 1991 Conversion of GDP into GTP by nucleoside diphosphate kinase on the GTP-binding proteins. *J Biol Chem* 256:21536-21540
518. Randazzo PA, Northrop JK, Kahn RA 1991 Activation of a small GTP-binding protein by nucleoside diphosphate kinase. *Science* 254:850-853
519. Donaldson JG, Finazzi D, Klausner RD 1992 Brefeldin A inhibits Golgi membrane-catalyzed exchange of guanine nucleotide onto ARF protein. *Nature* 360:350-352
520. Helms JB, Rothman JE 1992 Inhibition by brefeldin A of a Golgi membrane enzyme that catalyzes exchange of guanine nucleotide bound to ARF. *Nature* 360:352-354
521. Okabe-Kado J, Kasukabe T, Honma Y, Hayashi M, Henzel WJ, Hozumi M 1992 Identity of a differentiation inhibiting factor for mouse myeloid leukemia cells with nm23/nucleoside diphosphate kinase. *Biochem Biophys Res Commun* 182:987-994
522. Postel E, Berberich S, Flint S, Ferrone C 1993 Human c-myc transcription factor PuF identified as nm23-H2 nucleoside diphosphate kinase, a candidate suppressor of tumor metastasis. *Science* 261:478-480

TE

De
MeI.
II.
III.

IV.

V.
VI.

A
ity
inte
enc
cha:
ami
ple
tere
t on
stim
ove:
pop
that
peop
year
heal
the
bilit
Ti
age-
latio
indr

—
At
of Ep
60 C
the S